




New York State Department of Health



2003-2004 Health Advisories

Chemicals in Sportfish and Game



These advisories are also available from the New York State Department of Health Web site on the Internet: <http://www.health.state.ny.us/nysdoh/environ/fish.htm>

In an effort to reduce the costs of printing, please notify us if you wish your name to be deleted from our mailing list or if your address has changed. Comments regarding the format or content of this booklet are welcome. Use the telephone number for Environmental Health Information listed on page 23 or e-mail: BTSA@health.state.ny.us

2003-2004 Health Advisories: Chemicals in Sportfish and Game

Summary

The New York State Department of Health (DOH) issues advisories on eating sportfish and game because some of these foods contain chemicals at levels that may be harmful to your health. These advisories are for sportfish and game that people take and are not for fish and game sold in markets. The health advisories are: (1) general advice on sportfish taken from waters in New York State; (2) advice on sportfish from specific waterbodies; and (3) advice on eating game. The advisory tells you how to minimize your exposure to contaminants in sportfish and game and reduce whatever health risks are associated with them. The advisories are updated yearly.

Background

Fish and game are nutritious and good to eat. But some fish may take in contaminants from the water they live in and the food they eat. Game, too, may take in contaminants from their food and water. Some of these contaminants build up in fish and game--and in people--over time. These contaminants could harm people, so it is important to keep your exposure to these contaminants as low as possible.

The federal government sets standards for chemicals in food that is sold commercially, including fish. The decision to eat sportfish or game that you take is not regulated by government. Instead, state governments issue advisories. In New York State, the Department of Environmental Conservation (DEC) routinely monitors contaminant levels in fish and game and DOH issues advisories when sportfish have contaminant levels greater than federal standards.

These advisories are intended to allow you to select fish and game with lower contaminant levels so you can minimize your exposure to contaminants.

Health Benefits

When properly prepared, fish provide a diet high in protein and low in saturated fats. Almost any kind of fish may have real health benefits if it replaces a high-fat source of protein in the diet. You can get the health benefits of fish and reduce unwanted contaminants by following the guidelines in these advisories.

Contaminants in Fish and Game

Long-lasting contaminants, such as PCBs, DDT and cadmium, build up in your body over time. It may take months or years of regularly eating contaminated fish or game to build up amounts that are a health concern. Health problems that may result from the contaminants found in fish or game range from small changes in health that are hard to detect to birth defects and cancer. Mothers who eat highly contaminated fish and game before becoming pregnant may have children who are slower to develop and learn. The meal advice in this advisory is also intended to protect children from these potential developmental problems. Women beyond their childbearing years and men face fewer health risks from contaminants than children do. People in this group should follow the advisory to reduce their total exposure to contaminants.

Some contaminants cause cancer in animals. We cannot predict with certainty your risks of cancer from eating contaminated fish or game. Cancer currently affects about one in every three people, primarily due to smoking, diet and hereditary risk factors. Exposure to contaminants in the fish and game you eat may not increase your cancer risk at all. If you follow this advisory over your lifetime, you will minimize your exposure and reduce whatever cancer risk is associated with these contaminants.

More information about the chemicals that

have led to advisories in New York State sportfish and game and potential health effects can be found on page 19. When the federal government sets standards for fish, it generally assumes that people eat about a half-pound of fish each month. The contaminant levels are measured in a skin-on fillet which has not been trimmed; this sample is used in determining whether or not the fish exceeds standards. Fish cannot be legally sold if they contain a contaminant at a level greater than its standard. When sportfish from a waterbody contain contaminants at levels greater than the federal standards, DOH issues a specific advisory.

General Advisory for Eating Sportfish

The general health advisory for sportfish is that you eat no more than one meal (one-half pound) per week of fish taken from the state's freshwaters and some marine waters at the mouth of the Hudson River. These include the New York waters of the Hudson River, Upper Bay of New York Harbor (north of Verrazano Narrows Bridge), Arthur Kill, Kill Van Kull, Harlem River and the East River to the Throgs Neck Bridge (see map on page 17). This general advisory is to protect against eating large amounts of fish that have not been tested or may contain unidentified contaminants. The general advisory does not apply to most marine waters.

Specific Advisories for Freshwater, the Hudson River and the Upper Bay of New York Harbor

Fish from more than 80 waterbodies in New York have contaminant levels that are greater than federal standards. For these waters, DOH recommends either limiting or not eating a specific kind of fish (see pages 7 to 15). In some cases, enough information is available to issue advisories based on the length of the fish. Older (larger) fish are often more contaminated than younger (smaller) fish.

The contaminants that led to the advisory (mercury, cadmium, PCBs, chlordane, dioxin, DDT and mirex) are listed next to each advisory. If you eat fish from more than one water body with these advisories, you should limit consumption from all of the waters you

fish. For example, if you eat a meal of Koppers Pond carp, you should not eat American eel from Kinderhook Lake for the rest of that month since both of these fish species have EAT NO MORE THAN ONE MEAL PER MONTH advisories and both are based on PCB contamination.

Advisory for Women, Infants and Children

Health advice is also given for infants, children under the age of 15 and women of childbearing age. DOH recommends that these groups not eat any fish from the specific waterbodies listed in the advisory. The reason for this specific advice is that chemicals may have a greater effect on developing organs in young children or in the fetus. They also build up in women's bodies and are often passed on in mother's milk. Waters that have specific advisories have at least one species of fish with an elevated contaminant level, which means that a contamination source is or was in or near the water.

When eating fish from waters where cadmium or mercury are listed as primary contaminants, it is important to space out fish meals according to the specific advisory for that waterbody. For example, if you eat a meal of yellow perch from Moshier Reservoir, you should not eat any more fish with the same mercury advisory for the rest of that month. However, for other contaminants, the total number of meals that you eat during the year is important and many of those meals can be eaten during a few months of the year. If most of the fish you eat are from the ONE MEAL PER WEEK category, you should not exceed 52 meals per year. Likewise, if most of the fish you eat are in the ONE MEAL PER MONTH category, you should not exceed 12 meals per year. Remember, eating one meal of fish from the ONE MEAL PER MONTH group is the same as eating four meals from the ONE MEAL PER WEEK group.

U.S. Food and Drug Administration Advice

Due to concerns about mercury contamination, the U.S. Food and Drug Administration (FDA) advises pregnant women, women of childbearing age who may become

pregnant, nursing mothers and young children to eat no shark, swordfish, king mackerel or tilefish. FDA also acknowledges seafood can be an important part of a balanced diet for pregnant women and women who may become pregnant, and advises that they can eat up to 12 ounces per week of a variety of other kinds of fish. If you have any questions or would like additional information on this advice, you can call FDA (toll-free) at 1-888-INFO-FDA.

The full FDA advisory can be found at: <http://www.cfsan.fda.gov/~dms/admehg.html>. Additional information for consumers about mercury in fish from the market can be found at <http://www.cfsan.fda.gov/~frf/sea-mehg.html>.

Advisories for Other Marine Waters

DOH also issues specific advisories for Long Island Sound, Block Island Sound, Peconic/Gardiners Bays, the Lower Bay of New York Harbor, Jamaica Bay and other Long Island south shore waters (see maps on pages 17 and 18). These apply to striped bass, bluefish and American eels and are the only fish advisories that apply to these waters. Ocean fish, although tested less often, are generally less contaminated than freshwater fish. However, striped bass, bluefish and eels have specific habits or characteristics that make them more likely to have contaminants than other marine species (see page 18).

Advisories for Chemical Contaminants in Crabs and Lobsters, Including Hepatopancreas Advice

DOH strongly recommends that you not eat the soft green substance (mustard, tomalley, liver or hepatopancreas) found in the body section of crabs and lobsters from any waters, because cadmium, PCBs and other contaminants concentrate there. Because contaminants in this substance are transferred to cooking liquid, you should also discard crab or lobster cooking liquid. DOH also has a special advisory to eat no more than six Hudson River blue crabs per week.


Advisories for Eating Game

DOH also issues advisories about eating some game. These are on page 18 of this

booklet and include advisories for eating snapping turtles and waterfowl statewide because they contain PCBs and other contaminants. Because these contaminants concentrate in fat, you can minimize your exposure by not eating fat from these game and by following the cooking and eating advice on page 18.

Cleaning and Cooking Your Fish

Many contaminants are found at higher levels in the fat of fish. You can reduce the amount of these contaminants in a fish meal by properly trimming, skinning and cooking your catch. Remove the skin and trim all the fat from the belly flap, the line along the sides, the fat along the back and under the skin (see the diagram below).



Cooking or soaking fish cannot eliminate the contaminants, but heat from cooking melts some of the fat in fish and allows some of the contaminated fat to drip away. Broil, grill or bake the trimmed, skinned fish on a rack so that the fat drips away. Do not use drippings to prepare sauces or gravies.

These precautions will not reduce the amount of mercury or other metals. Mercury is distributed throughout a fish's muscle tissue (the part you eat), rather than in the fat and skin. Therefore, the only way to reduce mercury intake is to reduce the amount of contaminated fish you eat.

Reducing Exposure To Chemical Contaminants From Fish

Fish are an important source of protein and are low in saturated fat. Naturally occurring fish oils lower plasma cholesterol and triglycerides, thereby decreasing the risk of coronary heart disease. Increasing fish consumption is useful in reducing dietary fat and controlling weight. By eating a diet that includes food from a variety of protein sources, an individual is more likely to have a diet that is adequate in all nutrients.

Although eating fish has some health benefits, fish with high contaminant levels should be avoided. When deciding whether or not to eat fish that may be contaminated, the benefits of eating those fish can be weighed against the risks.

For young women, eating contaminated fish is a health concern not only for themselves but also for any unborn or nursing child, since the chemicals may reach the unborn babies and can be passed on in mother's milk. For an older person with heart disease, the risks, especially of long-term health effects, may not be as great a concern when compared to the benefits of reducing the risks of heart disease.

Everyone can benefit from eating the fish they catch and can minimize their contaminant intake by following these general recommendations:

1. Choose sportfish from waterbodies that are not listed on pages 7 through 15 and follow the advice in this booklet.
2. When preparing sportfish, use a method of filleting the fish that will reduce the skin, fatty material and dark meat. These parts of the fish contain many of the contaminants.
3. When deciding which sportfish to eat, choose smaller fish, consistent with DEC regulations, within a species since they may have lower contaminant levels. Older (larger) fish within a species may be more contaminated because they have had more time to accumulate contaminants in their bodies.
4. Do not eat the soft green tissue (mustard, tomalley, liver or hepatopancreas) found in the body section of crab and lobster. This tissue has been found to contain high levels of chemical contaminants, including PCBs and heavy metals.
5. When eating sportfish, use cooking methods such as broiling, poaching, boiling and baking, which allow contaminants from the fatty portions of fish to drain out. Pan frying is not recommended. The cooking liquids and fat drippings of fish from contaminated waters should be discarded since these liquids may retain contaminants.
6. Anglers who want to enjoy the fun of fishing but who wish to eliminate the potential risks associated with eating contaminated sportfish may want to consider "catch and release" fishing. Refer to the DEC *New York State Fishing Regulations Guide* for suggestions on catch and release fishing techniques.

Lead in Fishing Tackle and Bullets

Lead can cause health problems when it builds up in the body. Because the fetus and young child are at the greatest risk, it is particularly important for pregnant women, women of child-bearing age and young children to minimize their lead exposures. Lead poisoning can slow a child's physical growth and mental development and can cause behavior and other nervous system problems, reproductive problems, kidney and liver damage, blindness and even death in both adults and children. Fishing tackle (especially sinkers and jig heads), bullets and shot often contain lead; in order to reduce exposure to the lead in these products, you should:

- Keep all lead objects away from young children (young children often put their hands and objects in their mouth).
- Wash hands with soap and water after holding or using lead sinkers and jig heads or reloading lead bullets or shot.
- Never put lead sinkers in your mouth. This includes biting down on lead sinkers.
- Never eat, drink, or smoke immediately after handling lead sinkers, bullets or shot; wash hands first.
- Take proper precautions when melting lead and pouring sinkers or bullets at home. Use jacketed bullets (a jacketed bullet is a bullet enveloped in a casing of another material such as copper) and shotgun shells with plastic wads. Studies have shown that people can be exposed to

lead from shooting at indoor and outdoor firing ranges. For additional information on how to minimize your exposure to lead, call 1-800-458-1158, ext 27900 (toll free).

- Consider non-lead alternatives. The New York State Department of Environmental Conservation (DEC) encourages anglers to use non-lead alternatives for sinkers and jig heads to reduce the risk of lead poisoning to birds. More information is provided on the DEC website at: [http://www.dec.state.ny.us/website/dfwmr/fish/respangl.html#non toxic](http://www.dec.state.ny.us/website/dfwmr/fish/respangl.html#non%20toxic)

Good Sanitary Practices - Bacteria, Viruses and Parasites in Fish and Game

Fish and game and other meats can be contaminated with bacteria, viruses or parasites that can cause illness. You should harvest fish and game that act and look healthy, and follow good sanitary practices when preparing them. We recommend that you wear rubber or plastic protective gloves while filleting, field dressing, skinning or butchering. We also recommend that you remove intestines soon after harvest, don't eat intestines and avoid direct contact with intestinal contents. Hands, utensils and work surfaces should be washed before and after handling any raw food, including fish and game meat. Fish and game should be kept cool (with ice or refrigerated below 45EF or 7EC) until filleted or butchered and then should be refrigerated or frozen. Some hunters prefer to hang big game for several days before butchering; this should not be done unless the game can be kept at temperatures consistently below 45EF. Fish and other seafood should be cooked to an internal temperature (in the thickest part) of 140EF (60EC); game birds and other types of wild game meat should be cooked to an internal temperature (in the thickest part) of 165EF (74EC).

Advice on Eating Raw or Partially Cooked Fish, Shellfish and Other Meats

Foods of animal origin, such as pork, poultry, beef, dairy products, fish and shellfish, can be contaminated with bacteria, viruses or parasites that can cause illness. Persons at high risk (for example, those who are immunocompromised, suffer from liver disease or other chronic diseases) can be more susceptible to and more severely affected by these infectious diseases. This is why the

Department of Health recommends that all of these foods be thoroughly cooked before eating. Government agencies, universities and the food industry have active programs that strive to minimize contamination of raw animal foods and assure safe food products.

Information on rules and regulations, including areas in which clam, oyster and mussel collection is permitted, can be obtained from DEC by calling (631) 444-0475. DEC routinely tests clam, oyster and mussel beds for bacteria. Based on these tests, an area may be closed to clam, oyster and mussel harvesting. You can check the DEC website at <http://www.dec.state.ny.us/website/dfwmr/marine/shellfish/index.html> for general information on shellfish harvest. Call DEC at (631) 444-0480 for the latest information on emergency closures.

Deformed or Abnormal Fish

The health implications of eating deformed or abnormal fish are unknown. Any obviously diseased fish (marked by tumors, lesions or other abnormal condition of the fish skin, meat or internal organs) should be discarded.

Botulism in Fish and Waterfowl

In recent years, large numbers of some species of Lake Erie fish and waterfowl have been found dead, sick and dying, many of them as a result of botulism poisoning. The botulism poison is produced by *Clostridium botulinum*, a bacterium that is common in the environment and can produce harmful levels of botulism poison under certain environmental conditions. This poison has been found in some of the affected fish and waterfowl. The botulism poison can cause illness and death if consumed by humans or animals. Cooking may not destroy the botulism poison. This problem may also occur in other waters, and we don't know whether all or only some fish and waterfowl species can be affected.

No human cases of botulism poisoning have been linked to these events. However, as a precaution, do not eat any fish or game if they are found dead or dying, act abnormally or seem sick. If you must handle dead or dying fish, birds or other animals, cover your hands with disposable rubber or plastic protective gloves or a plastic bag. The New York State Department of Environmental Conservation is monitoring the situation and investigating the

cause of this problem.

Rabies and Chronic Wasting Disease (CWD) in Deer

Rabies and Chronic Wasting Disease (CWD) are two diseases that can cause abnormal behavior in deer. Rabies can be found in any mammal (especially raccoons, bats, skunks and foxes) and is found only occasionally in New York State deer. Chronic Wasting Disease (CWD) is a disease of deer and elk. CWD has not been found in deer taken in New York State but may be present in some deer or elk from several Western and mid-Western states and some Canadian provinces.

Rabies is a viral infection which causes a rapidly progressive disease of the animal's nervous system that leads to paralysis and always death, usually within several days after signs of the disease first appear. Rabid deer may seem to lose their normal fear of humans, appear to have injured hind legs, salivate excessively, or be found laying on the ground struggling. Rabies can be transmitted from infected mammals to humans by exposure to infected tissues, particularly nervous tissue and saliva. Treatment is available to prevent development of rabies in exposed humans. Rabies is almost always fatal in exposed humans who develop the disease. Thorough cooking will inactivate the rabies virus (see "Good Sanitary Practices" on page 5 of this booklet). Hunters should be aware that deer with rabies may have symptoms similar to CWD.

CWD is a brain infection of deer and elk that leads to loss of body functions, poor body condition and abnormal behavior such as staggering or very poor posture. It eventually leads to the death of the animal. CWD appears to be caused by abnormal, infectious proteins called prions. There is currently no evidence suggesting CWD causes disease in people; however, researchers are still studying this topic. Cooking does not destroy CWD.

The following precautions are recommended to lessen disease risk:

- Do not handle or eat any deer, or other game that appear sick, act strangely, or are found dead.
- Wear rubber or latex gloves when field dressing game.

- Use separate dedicated knives, saws and cutting boards to butcher deer and remove antlers, particularly when cutting through the skull or spinal cord. Do not use regular kitchen utensils.
- Wash thoroughly with soap and water any knives, butchering tools, work surfaces, hands and any other body parts that have been exposed to animal tissue, blood, urine or feces.
- Avoid handling the brain and spinal tissues or fluids, saliva and mouthparts of game animals. If these tissues or fluids are handled, wash hands thoroughly with soap and water. If these tissues or fluids make contact with a person's eyes, nose, mouth, or fresh open breaks in a person's skin, contact the local health department to see if rabies exposure may have occurred and whether the animal should be tested for rabies.
- If possible, request that the meat from your deer be processed separately, without adding other hunters' deer meat.

In general, the brain, spinal cord, liver, spleen, kidney, heart, eyes, tonsils, bone marrow and lymph nodes of game may pose a greater risk of infection for a number of diseases. Normal field dressing will eliminate most of these organs and tissues. Lymph nodes and bone marrow can be eliminated by boning out the meat and carefully trimming the fat and connective tissue. You may want to consider these precautions with New York State deer. However, these precautions should definitely be followed with deer and elk from states and Canadian provinces with confirmed CWD.

For additional information about handling, processing, or eating deer or elk meat in other states, contact the state agriculture, wildlife or health agencies in those states.

For additional information about CWD and rabies, visit the DOH web page at <http://www.health.state.ny.us/nysdoh/zoonoses/zoonoses.htm>, or call DOH (see page 23 for toll-free number).

For an update on CWD testing in New York State, visit the DEC website at <http://www.dec.state.ny.us/website/dfwmr/wildlife/deer/cwd.html>.

2003-2004 Health Advisories

The following recommendations are based on contaminant levels in fish and game. To minimize potential adverse health impacts, the DOH recommends:

- C **Eat no more than one meal (one-half pound) per week** of fish from the state's freshwaters, the Hudson River estuary, Upper Bay of New York Harbor (north of the Verrazano Narrows Bridge), Arthur Kill, Kill Van Kull, East River to the Throgs Neck Bridge and Harlem River, except as recommended below.
- C **Women of childbearing age, infants and children under the age of 15 should not eat** any fish species from waters listed below.
- C **Follow trimming and cooking advice.**
- C Observe the following restrictions on eating fish from these waters and their tributaries to the first barrier impassable by fish.
- C Advice for other marine waters is on page 18.

Water (County)	Species	Recommendations	Chemical(s) of Concern
<u>Amawalk Reservoir</u> [64] (Westchester)	Largemouth and smallmouth bass over 16"	Eat no more than one meal per month	Mercury
<u>Arthur Kill</u> [71] (Richmond)	See Hudson River (south of Catskill)		
<u>Ashokan Reservoir</u> [58] (Ulster)	Smallmouth bass over 16" and walleye	Eat no more than one meal per month	Mercury
<u>Barge Canal</u> [4] Tonawanda Creek from Lockport to Niagara River (Erie & Niagara)	Carp	Eat no more than one meal per month	PCBs
<u>Beaver Lake</u> [35] (Lewis)	Chain pickerel	Eat no more than one meal per month	Mercury
<u>Belmont Lake</u> [83] (Suffolk)	Carp	Eat no more than one meal per month	Chlordane, PCBs
<u>Big Moose Lake</u> [30] (Herkimer)	Yellow perch over 9 inches	Eat no more than one meal per month	Mercury
<u>Bog Brook Reservoir</u> [62] (Putnam)	Walleye over 21 inches	Eat no more than one meal per month	Mercury

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Boys Corner Reservoir [59] (Putnam)	Largemouth bass over 16 inches and walleye	Eat no more than one meal per month	Mercury
Buffalo River/Harbor [6] (Erie)	Carp	Eat none	PCBs
Canadice Lake [9] (Ontario)	Lake or brown trout	Eat no more than one meal per month	PCBs
Canandaigua Lake [10] (Ontario & Yates)	Lake trout over 24"	Eat no more than one meal per month	PCBs
<u>Cannonsville Reservoir</u> [53] (Delaware)	Smallmouth bass over 15" and yellow perch	Eat no more than one meal per month	Mercury
Carry Falls Reservoir [21] (St. Lawrence)	Walleye	Eat no more than one meal per month	Mercury
Cayuga Creek [2] (Niagara)	All species	Eat none	Dioxin
Chenango River [49]	Walleye over 22"	Eat no more than one meal per month	Mercury
Cranberry Lake [25] (St. Lawrence)	Smallmouth bass	Eat no more than one meal per month	Mercury
Cross River Reservoir [66] (Westchester)	Largemouth and smallmouth bass over 16"	Eat no more than one meal per month	Mercury
Dart Lake [31] (Herkimer)	Yellow perch over 10"	Eat no more than one meal per month	Mercury
Delaware Park Lake [5] (Erie)	Carp	Eat no more than one meal per month	PCBs
<u>Diverting Reservoir</u> [61] (Putnam)	Walleye	Eat no more than one meal per month	Mercury
<u>East Branch Reservoir</u> [63] (Putnam)	Walleye	Eat no more than one meal per month	Mercury
East River [70] (NYC)	American eel	Eat none	PCBs
	Atlantic needlefish, bluefish, striped bass and white perch	Eat no more than one meal per month	PCBs

Waters with changes from the 2002-2003 Health Advisories are **underlined**.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Eighteen Mile Creek [3] (Niagara)	All species	Eat none	PCBs
Ferris Lake [41] (Hamilton)	Yellow perch over 12"	Eat none	Mercury
	Smaller yellow perch	Eat no more than one meal per month	Mercury
Fourth Lake [37] (Herkimer & Hamilton)	Lake trout	Eat none	DDT
Francis Lake [33] (Lewis)	Yellow perch	Eat no more than one meal per month	Mercury
Freeport Reservoir [81] (Nassau)	Carp	Eat no more than one meal per month	Chlordane
Grant Park Pond [76] (Nassau)	Carp	Eat no more than one meal per month	PCBs
Grasse River [20] Mouth to Massena Power Canal (St. Lawrence)	All species	Eat none	PCBs
Halfmoon Lake [34] (Lewis)	Yellow perch	Eat no more than one meal per month	Mercury
Hall's Pond [77] (Nassau)	Carp and goldfish	Eat none	Chlordane
Harlem River [69] (NYC)	American eel	Eat none	PCBs
	Atlantic needlefish, bluefish, striped bass and white perch	Eat no more than one meal per month	PCBs
Herrick Hollow Creek [51] (Delaware)	Brook trout	Eat no more than one meal per month	PCBs
Hoosic River [44] (Rensselaer)	Brown trout over 14"	Eat no more than one meal per month	PCBs
Hudson River [48] Corinth Dam to Dam at Route 9 Bridge in South Glens Falls	Smallmouth bass over 14"	Eat no more than one meal per month	Mercury

Hudson River Advisory continued on next page

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
<u>Hudson River (continued)</u> Sherman Island Dam downstream to Feeder Dam at South Glens Falls	Carp	Eat no more than one meal per month	PCBs
Dam at Route 9 Bridge in South Glens Falls to Troy Dam	All species	Eat none	PCBs
Troy Dam south to bridge at Catskill	All species except Alewife, American shad, blueback herring, rock bass and yellow perch	Eat none	PCBs
	Alewife, blueback herring, rock bass and yellow perch	Eat no more than one meal per month	PCBs
	American shad (general advisory)	Eat no more than one meal per week	PCBs
Bridge at Catskill south to and including the Upper Bay of New York Harbor (north of Verrazano Narrows Bridge), Arthur Kill and Kill Van Kull	Gizzard shad	Eat none	PCBs
	American eel, Atlantic needlefish, bluefish, brown bullhead, carp, channel catfish, goldfish, largemouth bass, smallmouth bass, rainbow smelt, striped bass, walleye, white catfish and white perch	Eat no more than one meal per month	PCBs
	Blue crab	Eat no more than six crabs per week	Cadmium, PCBs
	--hepatopancreas	Eat none	Cadmium, PCBs
	--cooking liquid	Discard	Cadmium, PCBs
Dobbs Ferry south to Greystone	American eel	Eat none	PCBs
	Other species	See advisories for Hudson River south of Catskill (above)	

Waters with changes from the 2002-2003 Health Advisories are **underlined**.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Indian Lake [17] (Lewis)	All species	Eat no more than one meal per month	Mercury
Irondequoit Bay [8] (Monroe)	Carp	Eat none	PCBs, Mirex
Keuka Lake [11] (Yates & Steuben)	Lake trout over 25"	Eat no more than one meal per month	DDT
Kill Van Kull [72] (Richmond)	See Hudson River (south of Catskill)		PCBs
Kinderhook Lake [47] (Columbia)	American eel	Eat no more than one meal per month	PCBs
Koppers Pond [12] (Chemung)	Carp	Eat no more than one meal per month	PCBs
Lake Capri [84] (Suffolk)	American eel and carp	Eat no more than one meal per month	Chlordane, Cadmium
Lake Champlain [23] Whole Lake	Lake trout over 25" and walleye over 19"	Eat no more than one meal per month	PCBs, Mercury
Bay within Cumberland Head to Crab Island	Brown bullhead	Eat none	PCBs
	American eel and yellow perch	Eat no more than one meal per month	PCBs
Lake Ontario [7] Including Niagara River below Niagara Falls (see Niagara River for additional advice)	American eel, channel catfish, carp, lake trout over 25", brown trout over 20" and chinook salmon	Eat none	PCBs, Mirex, Dioxin
	White sucker, rainbow trout, smaller lake trout, smaller brown trout and coho salmon over 25"	Eat no more than one meal per month	PCBs, Mirex, Dioxin
West of Point Breeze	White perch	Eat none	PCBs, Mirex, Dioxin
East of Point Breeze	White perch	Eat no more than one meal per month	PCBs, Mirex, Dioxin
Loft's Pond [79] (Nassau)	Carp and goldfish	Eat no more than one meal per month	Chlordane

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Long Pond-Croghan [26] (Lewis)	Splake over 12"	Eat none	Mercury
Lower and Upper Sister Lakes [29] (Hamilton)	Yellow perch over 10"	Eat none	Mercury
Upper Massapequa Reservoir [82] (Nassau)	White perch	Eat no more than one meal per month	Chlordane
Massena Power Canal [19] (St. Lawrence)	Smallmouth bass	Eat no more than one meal per month	PCBs
Meacham Lake [22] (Franklin)	Yellow perch over 12"	Eat none	Mercury
	Smaller yellow perch	Eat no more than one meal per month	Mercury
Mohawk River [42] Between Oriskany and West Canada Creeks (Oneida & Herkimer)	Carp	Eat none	PCBs
	Largemouth bass and tiger muskellunge	Eat no more than one meal per month	PCBs
Moshier Reservoir [27] (Herkimer)	Yellow perch	Eat no more than one meal per month	Mercury
Nassau Lake [46] (Rensselaer)	All species	Eat none	PCBs
Neversink Reservoir [56] (Sullivan)	Smallmouth bass	Eat no more than one meal per month	Mercury
New York Harbor [73]	See Hudson River (south of Catskill) and marine waters advice on page 17		PCBs
Niagara River [1] Above Niagara Falls	Carp	Eat no more than one meal per month	PCBs
	White perch	Eat none	PCBs, Mirex, Dioxin
Below Niagara Falls (also see Lake Ontario)	Smallmouth bass	Eat no more than one meal per month	PCBs, Mirex, Dioxin

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Onondaga Lake [14] (Onondaga)	Walleye	Eat none	Mercury
	Carp and Channel catfish	Eat no more than one meal per month	Dioxin, PCBs, Mercury
	All other species	Eat no more than one meal per month	Mercury
Oswego River [15] Oswego power dam to upper dam at Fulton (Oswego)	Channel catfish	Eat no more than one meal per month	PCBs
Pepacton Reservoir [55] (Delaware)	Smallmouth bass over 15"	Eat no more than one meal per month	Mercury
Ridders Pond [75] (Nassau)	Goldfish	Eat none	Chlordane
Rondout Reservoir [57] (Sullivan and Ulster)	Smallmouth bass over 16"	Eat no more than one meal per month	Mercury
Round Pond [38] Town of Long Lake (Hamilton)	Yellow perch over 12"	Eat no more than one meal per month	Mercury
St. James Pond [85] (Suffolk)	All species	Eat no more than one meal per month	Chlordane, DDT
St. Lawrence River [18] Whole River	American eel, channel catfish, lake trout over 25", carp, brown trout over 20" and chinook salmon	Eat none	PCBs, Mirex, Dioxin
	White perch, white sucker, rainbow trout, smaller lake trout, smaller brown trout and coho salmon over 25"	Eat no more than one meal per month	PCBs, Mirex, Dioxin
Bay at St. Lawrence/ Franklin Co. line	All species	Eat none	PCBs
Salmon River [16] Mouth to Salmon Reservoir (Oswego) (also see Lake Ontario)	Smallmouth bass	Eat no more than one meal per month	PCBs, Mirex

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.

Water (County)	Species	Recommendations	Chemical(s) of Concern
Sauquoit Creek [43] Between Old Silk Mill Dam (near New Hartford/Paris town line) and Mohawk River (Oneida)	Brown trout	Eat none	PCBs
Saw Mill River [67] (Westchester)	American eel	Eat no more than one meal per month	Chlordane
Schoharie Reservoir [54] (Delaware, Greene and Schoharie)	Walleye	Eat no more than one meal per month	Mercury
Schroon Lake [39] (Warren & Essex)	Lake trout over 27", yellow perch over 13" and smallmouth bass	Eat no more than one meal per month	PCBs, Mercury
Sheldrake River [68] (Westchester)	American eel	Eat none	Chlordane, Dieldrin
	Goldfish	Eat no more than one meal per month	Chlordane
Skaneateles Creek [13] From dam at Skaneateles to Seneca River (Onondaga)	Brown trout over 10"	Eat no more than one meal per month	PCBs
Smith Pond - Rockville Centre [78] (Nassau)	White perch	Eat no more than one meal per month	Chlordane
Smith Pond - Roosevelt Park [80] (Nassau)	American eel	Eat none	Chlordane
	Carp and goldfish	Eat no more than one meal per month	Chlordane
Soft Maple Reservoir [36] (Lewis)	Rock bass	Eat no more than one meal per month	Mercury
Spring Pond - Middle Island [86] (Suffolk)	Carp and goldfish	Eat none	Chlordane
Stillwater Reservoir [28] (Herkimer)	Yellow perch over 9", smallmouth bass and splake	Eat no more than one meal per month	Mercury

Waters with changes from the 2002-2003 Health Advisories are underlined.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.


Water (County)	Species	Recommendations	Chemical(s) of Concern
Sunday Lake [32] (Herkimer)	Yellow perch	Eat no more than one meal per month	Mercury
Susquehanna River [52]	Walleye over 22"	Eat no more than one meal per month	Mercury
Threemile Creek [40] (Oneida)	White sucker	Eat no more than one meal per month	PCBs
Titicus Reservoir [65] (Westchester)	White perch	Eat no more than one meal per month	Mercury
Tupper Lake [24] (Franklin & St. Lawrence)	Walleye	Eat no more than one meal per month	Mercury
Unadilla River [50]	Walleye over 22"	Eat no more than one meal per month	Mercury
Upper and Lower Sister Lakes [29] (Hamilton)	Yellow perch over 10"	Eat none	Mercury
Valatie Kill [45] Between County Rt. 18 and Nassau Lake (Rensselaer)	All species	Eat none	PCBs
West Branch Reservoir [60] (Putnam)	Walleye	Eat no more than one meal per month	Mercury
Whitney Park Pond [74] (Nassau)	Carp and goldfish	Eat no more than one meal per month	Chlordane

Waters with changes from the 2002-2003 Health Advisories are **underlined**.

Numbers in brackets refer to map on page 16.

Please note the special advice for **women of childbearing age, infants and children under the age of 15** on page 2.


Waters with Specific Fish Consumption Advisories 2003



- | | | |
|----------------------------|---------------------------------|----------------------------------|
| 1 Niagara River | 29 Upper and Lower Sister Lakes | 58 Ashokan Res. |
| 2 Cayuga Creek | 30 Big Moose Lake | 59 Boyds Corner Res. |
| 3 Eighteen Mile Creek | 31 Dart Lake | 60 West Branch Res. |
| 4 Barge Canal | 32 Sunday Lake | 61 Diverting Res. |
| 5 Delaware Park Lake | 33 Francis Lake | 62 Bog Brook Res. |
| 6 Buffalo River and Harbor | 34 Halfmoon Lake | 63 East Branch Res. |
| 7 Lake Ontario | 35 Beaver Lake | 64 Amawalk Res. |
| 8 Irondequoit Bay | 36 Soft Maple Res. | 65 Titicus Res. |
| 9 Canadice Lake | 37 Fourth Lake | 66 Cross River Res. |
| 10 Canandaigua Lake | 38 Round Pond | 67 Saw Mill River |
| 11 Keuka Lake | 39 Schroon Lake | 68 Sheldrake River |
| 12 Koppers Pond | 40 Threemile Creek | 69 Harlem River |
| 13 Skaneateles Creek | 41 Ferris Lake | 70 East River |
| 14 Onondaga Lake | 42 Mohawk River | 71 Arthur Kill |
| 15 Oswego River | 43 Sauquoit Creek | 72 Kill Van Kull |
| 16 Salmon River | 44 Hoosic River | 73 New York Harbor |
| 17 Indian Lake | 45 Valatie Kill | 74 Whitney Park Pond |
| 18 St. Lawrence River | 46 Nassau Lake | 75 Ridders Pond |
| 19 Massena Power Canal | 47 Kinderhook Lake | 76 Grant Park Pond |
| 20 Grasse River | 48 Hudson River | 77 Hall's Pond |
| 21 Carry Falls Res. | 49 Chenango River | 78 Smith Pond (Rockville Centre) |
| 22 Meacham Lake | 50 Unadilla River | 79 Loft's Pond |
| 23 Lake Champlain | 51 Herrick Hollow Creek | 80 Smith Pond (Roosevelt Park) |
| 24 Tupper Lake | 52 Susquehanna River | 81 Freeport Res. |
| 25 Cranberry Lake | 53 Cannonsville Res. | 82 Upper Massapequa Res. |
| 26 Long Pond (Croghan) | 54 Schoharie Res. | 83 Belmont Lake |
| 27 Moshier Res. | 55 Pepacton Res. | 84 Lake Capri |
| 28 Stillwater Res. | 56 Neversink Res. | 85 St. James Pond |
| | 57 Rondout Res. | 86 Spring Pond (Middle Island) |

P:\Bureau\Fish_Game\Advmaps\03map1.wor

Map of New York City Harbor Region



Additional Advice

Advisories for Lake Erie - Due to PCB contamination, women of childbearing age, infants and children under the age of 15 are advised to eat no more than one meal per week of chinook salmon less than 19 inches, burbot, freshwater drum, lake whitefish, rock bass and yellow perch and to EAT NO MORE THAN ONE MEAL PER MONTH of all other fish from Lake Erie. Other people should eat no more than one meal per week of any Lake Erie fish species.

Marine Bluefish and Eels - The general advisory {Eat no more than one meal (one-half pound) per week} applies to bluefish and American eels but not to most other fish (see Marine Striped Bass below) from Long Island Sound, Block Island Sound, Peconic/Gardiners Bays, the Lower Bay of New York Harbor, Jamaica Bay and other Long Island south shore waters. (Contaminant of concern - PCBs)

Marine Striped Bass - Women of childbearing age and children under the age of 15 should eat no striped bass from Upper and Lower Bays of New York Harbor or Long Island Sound west of Wading River. Other people should EAT NO MORE THAN ONE MEAL PER MONTH of striped bass from these waters. Everyone should eat no more than one meal per week of striped bass taken from Jamaica Bay, Eastern Long Island Sound, Block Island Sound, Peconic/Gardiners Bay or Long Island south shore waters. (Contaminant of concern - PCBs)

Crabs and Lobsters - The hepatopancreas (sometimes called mustard, tomalley or liver) of crabs and lobsters should not be eaten because it has high contaminant levels. Because contaminants in the hepatopancreas are transferred to cooking


liquid, crab or lobster cooking liquid should also be discarded. (Contaminants of concern - PCBs, cadmium, dioxin).

Hudson River American Shad - The advisory for women of childbearing age, infants and children under 15 is EAT NONE for all fish from the lower Hudson River because of PCB contamination. American shad have lower PCB levels than other species. A few meals of Hudson River American shad meat and roe, especially using cooking and trimming methods to minimize PCB content, would not pose an unacceptable health risk for women of childbearing age and children, assuming this is their only significant exposure to PCBs.

Snapping Turtles - Snapping turtles retain contaminants in their fat, liver, eggs and, to a lesser extent, muscle. If you choose to consume snapping turtles, you can reduce your exposure by carefully trimming away all fat and discarding the fat, liver and eggs prior to cooking the meat or preparing soup. Women of childbearing age, infants and children under the age of 15 should AVOID EATING snapping turtles or soups made with their meat. (Contaminant of concern - PCBs).

Wild Waterfowl - Mergansers are the most heavily contaminated waterfowl species and should NOT BE EATEN. EAT NO MORE THAN TWO MEALS PER MONTH of other wild waterfowl; you should skin them and remove all fat before cooking, and discard stuffing after cooking. Wood ducks and Canada geese are less contaminated than other wild waterfowl species and diving ducks are more contaminated than dabbling ducks. (Contaminants of concern - PCBs, mirex, chlordane, DDT)

Map of New York Marine Waters



Information on Chemicals in Sportfish and Game

The following paragraphs give some basic information on chemicals in sportfish and game in New York State. Most of our knowledge of potential health effects comes from high dose animal studies or worker exposures. Chemicals that cause adverse health effects in humans and laboratory animals after high levels of exposure may increase the risk of adverse effects in humans exposed to lower levels for long periods of time. Following the suggestions in the advisory will minimize your exposure and any health risks from contaminants in fish.

Chlordane

Chlordane is a man-made pesticide that was used widely to control agricultural and home/garden pests until most uses were banned in the United States during the mid-1970s. In New York State, chlordane was used for the underground control of termites until that use was banned in 1985. Chlordane generally got into bodies of water after improper waste disposal or run-off from treated areas. Chlordane builds up in the fatty tissues of fish, birds and mammals. Since chlordane is present in the fatty tissues of fish, exposure to chlordane in fish can be reduced by certain cleaning and cooking practices (see page 3).

People exposed to large amounts of chlordane may have nervous system damage. Exposure to high levels of chlordane damages the nervous system and liver of laboratory animals. Some animals exposed before birth and while nursing developed behavioral effects later. Chlordane causes cancer in laboratory animals exposed to high levels over their lifetimes. Whether chlordane causes cancer in humans is unknown.

DDT (also includes DDE and DDD)

DDT is a man-made pesticide that was used widely to control insects on agricultural crops and biting insects, such as mosquitos and black flies. Its use was banned in New York in 1971 and throughout the United States in 1973. DDT generally got into bodies of water after improper waste disposal, direct spraying

of water bodies or run-off from treated areas.

In the environment, DDT breaks down into DDE and DDD, chemicals which are very similar to DDT. DDT builds up in the fatty tissues of fish, birds and mammals. Since DDT is present in the fatty tissues of fish, exposure to DDT in fish can be reduced by certain cleaning and cooking practices (see page 3).

People who accidentally ingested large amounts of DDT had effects on the nervous system that went away once the exposure stopped.. A study in humans showed that increasing concentrations of p,p'-DDE in human breast milk were associated with reductions in the duration of lactation. An additional study in humans found that as the DDE levels in the blood of pregnant women increased, the chances of having a pre-term baby also increased. Exposure of laboratory animals to high levels of DDT damages the liver and can cause reproductive, developmental and nervous system effects. DDT causes cancer in laboratory animals exposed to high levels over their lifetimes. Whether DDT causes cancer in humans is unknown.

Mirex

Mirex is a man-made chemical that was used as a pesticide to control fire ants until its use was banned in the United States in the late 1970s. It was also used as a flame retardant in plastics, rubber, paint, paper and electrical goods until the early 1970s. Mirex generally got into bodies of water after improper waste disposal or run-off from treated areas. Mirex builds up in the fatty tissues of fish, birds and mammals. Since mirex is present in the fatty tissues of fish, exposure to mirex in fish can be reduced by certain cleaning and cooking practices (see page 3).

Laboratory animals exposed to mirex had damage to the eyes, nervous system, reproductive system, liver, thyroid and kidneys. Mirex causes cancer in laboratory animals exposed to high levels over their lifetimes. Whether mirex causes cancer in humans is unknown.

PCBs

PCBs (polychlorinated biphenyls) are a family of man-made chemicals that were used in many commercial and electrical products until their manufacture was banned in the mid-1970s. Some electrical equipment still in use contains PCBs. In this country, most PCBs were sold as mixtures called Aroclors. PCBs build up in fatty tissues of fish, birds and mammals. Since PCBs are present in the fatty tissues of fish, exposure to PCBs in fish can be reduced by certain cleaning and cooking practices (see page 3). Industrial workers who were exposed to large amounts of PCBs and other chemicals in air (and perhaps through their skin) experienced skin, eye and respiratory tract irritation and mild changes in the functioning of their livers. Less frequently, workers exposed to high levels reported headache, digestive disturbances, and showed liver problems.

Some studies of pregnant women suggest a link between a mother's increased exposure to PCBs and other chemicals from eating contaminated fish or from other environmental sources and slight effects on her child's birthweight, short-term memory, and learning. Recent studies suggest that women who ate fish containing PCBs (and other contaminants) have slightly shorter menstrual cycles and take a longer time to become pregnant than women who did not eat contaminated fish. A study of older adults (49 to 86 years old) who ate fish containing PCBs (and other contaminants) suggests that PCB exposure is associated with lower scores on several measures of memory and learning. Although some of the studies did control for the possible effects of other chemical contaminants, the role of these chemicals in causing the observed effects is not fully understood.

Studies of workers exposed to PCBs in air (and perhaps through their skin) raise concerns about the human carcinogenicity of PCBs, but results are inconsistent. The data are inadequate to prove that PCB exposures cause cancer in humans.

PCBs affect the skin, liver, and the nervous, immune and reproductive systems of animals exposed to high levels. PCBs also reduce the birth weight and change the behavior of offspring born to animals exposed

before, during, and after pregnancy. A few individual PCBs cause birth defects in offspring born to animals exposed during pregnancy. Some types of PCBs cause cancer in laboratory animals exposed over their lifetime.

PCDDs and PCDFs

Polychlorinated dibenzo-p-dioxins (also known as PCDDs or dioxins) and chlorinated dibenzofurans (also known as PCDFs or furans) are two closely related families of chemical compounds. Some dioxins and furans are produced as unwanted by-products in chemical manufacturing processes, such as in the production of certain herbicides and disinfectants. They are also found in the smoke or ash from motor vehicles, municipal waste incinerators and wood fires. Some dioxins and furans are environmentally and biologically persistent. They are highly soluble in fats and are stored in the fatty tissue of fish and other animals. Since dioxins and furans are present in fatty tissues of fish, exposure to dioxins and furans in fish can be reduced by certain cleaning and cooking practices (see page 3).

Dioxins and furans are thought to produce similar health effects. TCDD (2,3,7,8-tetra-chlorodibenzo-p-dioxin) is the most potent of the dioxins and furans, and much of what we know about the toxicity of dioxins and furans comes from studies of TCDD.

Many studies have looked at how exposure to high level of dioxins and furans can affect human health. Most of these studies examined workers exposed during the manufacture of chemicals, including pesticides, contaminated with dioxins. Other studies have looked at people exposed during the use of pesticides contaminated with dioxins or at people exposed to dioxins and/or furans after an industrial accident.

The most obvious health effect in people exposed to large amounts of dioxins is chloracne (a severe acne-like skin disease). Other effects that are clearly caused by dioxins are other types of skin diseases and changes in the blood levels of reproductive hormones and of enzymes indicating changes in liver function. Other effects have been reported more than once, but the evidence for these effects is not

strong enough to clearly indicate that they were caused by dioxins. These include increases in cardiovascular disease, diabetes or altered glucose (sugar) metabolism, reproductive effects, nervous system effects, and thyroid function changes. Immunological and respiratory system effects also have been reported but results are only suggestive that these effects were caused by dioxins. Several studies showed a positive association between a person's risk of getting cancer (all type of cancers combined) and working in a plant where products contaminated with dioxins were manufactured, but the evidence is not strong enough to conclude the cancers were caused by dioxins and not some other factor or combination of factors.

Several studies have looked at the effects of dioxins and furans on the health of children. Children of Japanese and Taiwanese women who ingested rice oil highly contaminated with furans and other dioxin-like compounds showed developmental effects that ranged from very mild to very severe. Scientists in Europe have reported associations between a mother's exposures to dioxins and furans and effects on her child's mental and motor skills, immune system, play behavior, and hormone levels. However, the study has limitations and the effects may have been associated with other factors and not with the dioxins and furans.

In laboratory animals, TCDD has damaged the liver, skin, blood and the gastrointestinal, immune, reproductive and nervous systems. It also affects prenatal development in animals whose mothers were exposed to TCDD. TCDD causes cancer in animals exposed to high levels over their lifetime.

Mercury

Mercury is a metal that occurs naturally in the environment in several forms. The metallic, or elemental form is a silvery, odorless liquid which can evaporate at room temperature to form odorless, colorless mercury vapor. Most of the mercury that accumulates in the fleshy part of fish is in an organic (carbon-containing) form called methylmercury. Greater amounts of methylmercury are found in older fish that tend to eat other fish and organisms. Methylmercury is found throughout the part of the fish that is eaten; therefore cleaning and cooking methods

that may reduce exposure to other contaminants are NOT effective for reducing exposure to mercury.

Exposure to high levels of metallic, inorganic or organic mercury can damage the nervous system and kidneys. People who ate fish that contained large amounts of methylmercury had permanent damage to the brain, kidneys and fetus. Exposure to methylmercury is more of a concern for children and unborn babies because their nervous system is still developing and the nervous system is a target organ for mercury. To address this concern, two separate long-term studies are underway in the Seychelles and Faroe Islands to look for health effects among children in populations that eat large amounts of seafood containing low levels of methylmercury. In these studies, the children's health is assessed as they grow. To date, results from the two studies are contradictory. The Seychelles Islands study shows no adverse effects, while the Faroe Islands study shows an association of mercury exposure and problems with memory, attention and language development. Interpreting the Faroe Islands study results is complicated because that population is also exposed to PCBs from the marine mammals which they eat. Both studies are ongoing and will continue to evaluate the children's health as they grow.

Cadmium

Cadmium is a naturally occurring metal found at low levels in soil and water. Cadmium is used in many industrial operations and in consumer products such as paints, plastics and batteries. Food, air and drinking water all contribute to a person's exposure to cadmium. Cadmium can be found in food items and in tobacco. Vegetables, fruits and cereals are the greatest source of cadmium. Cadmium can also be found in fish and shellfish from waters containing cadmium.

Eating food or drinking beverages containing high levels of cadmium can cause nausea, vomiting, stomach upset, cramps and diarrhea. Because cadmium leaves the body slowly, it can accumulate in the body, mainly in the kidneys, with continuing exposure. Some people with long-term exposure had kidney, bone and blood damage.

Contacts for Additional Information

New York State Department of Health

For more information on **health effects** from exposure to chemical contaminants or to provide comments on the format or content of this report contact:

Environmental Health Information: 1-800-458-1158, extension 27815 (toll-free). Calls are taken from 8:00AM-4:30PM, Monday through Friday. After hours, leave a voice mail message. The full advisories are also available from the Internet: <http://www.health.state.ny.us/nysdoh/fish/fish.htm> or can be requested by e-mail: BTSA@health.state.ny.us

New York State Department of Environmental Conservation

For more information on **fishing inland waters**, contact:

Region 1

Loop Rd.
Bldg. 40 SUNY
Stony Brook, NY 11790
(631) 444-0280

Region 2

1 Hunter Point Plaza
47-40 21st St.
Long Island City, NY 11101-5407
(718) 482-4922

Region 3

21 South Putt Corners Rd.
New Paltz, NY 12561-1696
(845) 256-3161

Region 4

Rt. 10, Jefferson Rd.
Stamford, NY 12167-9503
(607) 652-7366

Region 5

Rt. 86, P.O. Box 296
Raybrook, NY 12977-0296
(518) 897-1333

Region 6

317 Washington St.
Watertown, NY 13601-3787
(315) 785-2262

Region 7

1285 Fisher Ave.
Cortland, NY 13045-1090
(607) 753-3095, ext.213

Region 8

6274 E. Avon-Lima Rd.
Avon, NY 14414-9519
(585) 226-5343

Region 9

270 Michigan Ave.
Buffalo, NY 14203-2999
(716) 851-7000 or 7010

For more information on **fishing marine waters**, contact:

Bureau of Finfish and Crustaceans
205 North Belle Mead Road, Suite 1
East Setauket, NY 11733
(631) 444-0435

For information on **contaminant levels**, in fish and shellfish and wildlife contact:

Bureau of Habitat
625 Broadway
Fifth Floor
Albany, NY 12233-4756
(518) 402-8996

Prepared by:
New York State Department of Health
Division of Environmental Health Assessment
May 27, 2003
P:\Bureau\Fish_Game\Advisory\03 fish.wpd



State of New York
George E. Pataki, Governor

Department of Health
Antonia C. Novello, M.D., M.P.H., Dr. P.H., Commissioner