













| | | | |
|-------------------|------------------------|--------------------|---------------|
| Otter Lake | Otter Lake Association | Town of Forestport | Oneida County |
|-------------------|------------------------|--------------------|---------------|

| | | | |
|---|----------------------------------|---|-----------|
|  | Lake Characteristics | Surface area (ac/ha) | 282 / 114 |
| | | Max depth (ft/m) | 10 / 3 |
| | | Mean depth (ft/m) | 5 / 1 |
| | | Retention time (years) | 0.4 |
| | | Lake Classification | A |
| | | Dam Classification | A |
| | Watershed Characteristics | Watershed area (ac /ha) | 1478/598 |
| | | Watershed / Lake ratio | 5 |
| | | Lake & wetlands % | 16% |
| | | Agricultural % | 0% |
| | | Forest, shrub, grasses % | 65% |
| | | Residential | 0% |
| CSLAP Participation | Years | 1992-1996, 2002-2009, 2011, 2013, 2015-2016 | |
| | Volunteers | Scott and Mona Lincoln | |

| | | | |
|----------------------|--|---|-----------------------|
| Trophic state | HABs Susceptibility | Invasive Vulnerability | PWL Assessment |
| Mesoeutrophic | No reported blooms, Low susceptibility | No invasives in area, Low Vulnerability | Stressed |


Water quality values for Otter Lake for the 2016 sampling season. "Seasonal change" shows current year variability. Light red color indicates eutrophic conditions in top table and bloom conditions in bottom table.

| Open Water Indicators | 2016 Sampling Results | | | | | | | | Seasonal change | Long Term Avg | Long Term Trend? |
|-----------------------|-----------------------|-------|-------|-------|-------|-------|-------|-------|---|---------------|------------------|
| | 6/6 | 6/14 | 6/26 | 7/10 | 7/25 | 8/5 | 8/28 | 9/12 | | | |
| Clarity (m) | 1.4 | 1.7 | 1.6 | 1.6 | 1.4 | 1.6 | 1.5 | 1.7 |  | 1.9 | ↑ |
| TP (mg/l) | 0.017 | 0.011 | 0.025 | 0.013 | 0.014 | 0.008 | 0.017 | 0.013 |  | 0.013 | no |
| Deep TP (mg/l) | | | | | | | | |  | | |
| TN (mg/l) | 0.500 | 0.354 | 0.436 | 0.603 | 0.333 | 0.421 | 0.505 | 0.411 |  | 0.480 | no |
| N:P Ratio | 29 | 32 | 18 | 45 | 25 | 51 | 30 | 31 |  | 37 | no |
| Chl.a (ug/l) | 1.3 | 5.3 | 10.1 | 4.8 | 3.1 | 3.6 | 5.1 | 5.2 |  | 5.3 | no |
| pH | 7.0 | 7.0 | 7.5 | 7.5 | 7.5 | 8.4 | 7.6 | 7.7 |  | 7.3 | no |
| Cond (umho/cm) | 77 | 94 | 116 | 110 | 104 | 85 | 104 | 112 |  | 106 | no |
| Upper Temp (degC) | 22 | 18 | 23 | 25 | 27 | 24 | 27 | 25 |  | 21 | no |
| Deep Temp (degC) | | | | | | | | |  | | |
| BG Chl.a (ug/l) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |  | 1 | no |
| HABs reported? | no | no | no | no | no | no | no | no | | | |




Shoreline bloom and HABs notifications

| | | | |
|--|----------------------|--------------------------------------|----------------------|
| Date of first listing | Date of last listing | # weeks on the DEC notification list | # Weeks with updates |
| | | | |
| Shoreline HAB sample dates 2016 | | | |
| HAB Indicators | HAB criteria | No shoreline HABs samples 2016 | |
| BGA | 25 - 30 ug/L | | |
| microcystin | 20 ug/L | | |
| anatoxin - a | 4 ug/L | | |

HABs Status Open water Algae




Shoreline Algae




Otter Lake Long Term Trend Analysis


Clarity




Chlorophyll a



Surface and Deep Phosphorus




Lake Perception




Otter Lake Long Term Trend Analysis


Nitrogen




pH



Temperature




Conductivity




Otter Lake In-Season Analysis






















In Season Temperature








In Season Water Clarity



Scorecard

| Lake Use | | | | |
|-------------------------|---|---|---|--------------------|
| | PWL | Average Year | 2016 | Primary issue |
| Potable Water |  |  |  | No impacts |
| Swimming |  |  |  | Decreasing clarity |
| Recreation |  |  |  | Decreasing clarity |
| Aquatic Life |  |  |  | No impacts |
| Aesthetics |  |  |  | Native plants |
| Habitat |  |  |  | No impacts |
| Fish Consumption |  |  |  | Not applicable |

 Supported / Good
 Threatened / Fair
 Stressed / Poor
 Impaired
 Not Known

Summary

2016 compared to prior years: Otter Lake continues to be mesoligotrophic, or moderately unproductive. Water clarity was slightly lower than usual in 2016, and water temperatures were higher. Each of the other indicators was close to normal.

Compared to nearby lakes: Otter Lake has slightly lower clarity compared to other nearby (western Adirondack region) lakes- this is probably due to shallower water. Aquatic plant coverage is similar to the plant coverage in many of these other lakes, despite the lack of any invasive species. Chloride levels are between the 25th and 50th percentile for New York state lakes, indicating a potential for aquatic life impacts from road salt.

Trends: Water clarity has decreased over the last two decades. Phosphorus and water temperatures appear to have increased, as algae levels have decreased, but these changes have not been statistically significant.

Algal blooms and HABS: Water quality conditions indicated a low susceptibility to blooms, and none have been reported on the lake. The algae community in the open water samples is comprised of several different taxa, particularly green algae, and overall algae levels are low.

Aquatic invasive species: No AIS have been documented in any waterbody within 10 miles of Otter Lake. This may be consistent with the lack of public access to the lake, and suggests a low vulnerability to AIS introductions.

Indicated Actions: Individual stewardship activities such as pumping your septic system, growing a buffer of native plants next to the water bodies, and reducing erosion from shoreline properties and runoff into the lake will help to improve lake health by reducing nutrient and sediment loading to the lake. Visiting boats should be inspected to reduce the risk of new invasive species, and continued monitoring for invasive species is warranted. Continued algae bloom education and monitoring for HABS is recommended.

How to Read the Report

Welcome to the new and improved Citizens Statewide Lake Assessment Program individual lake report! In order to make this individual lake report as easy to digest as possible for the average reader, the length of the report has been greatly reduced. We hope that presenting the data in a more succinct manner will draw in more readers and hold their attention. Unfortunately, this new format leaves little room for definitions of terms, so we are including this section primarily as a glossary of terms for which the average reader may not know the definition.

The report begins with the lake name, town, and county, as well as the current NYS Federation of Lake Associations association, if one exists. The next section contains some physical characteristics of the lake. The surface area is the two dimensional area of the lakes surface and is given in units of acres and hectares. The max depth is the water depth measured at the deepest part of the lake and is given in units of feet and meters. The mean depth is either known from a rigorous study of the bathymetry of the lake or is calculated as 0.46 times the maximum depth and is given in units of feet and meters. The retention time is the time it takes for a drop of water to pass through a lake, given in units of years. The lake classification is a letter defining the “best uses” for this particular lake, based on the legal classification assigned by New York state. Class AA, AAspec and A lakes may be used as sources of potable water. Class B lakes are suitable for contact recreational activities, like swimming. Class C lakes are suitable for non-contact recreational activities, including fishing, although they may still support swimming. The addition of a T or TS to any of these classes indicates the ability of a lake to support trout populations and/or trout spawning. The dam classification is a letter defining the hazard class of a dam if one exists. Class A, B, C, and D dams are defined as low, intermediate, high, or negligible/no hazard dams in that order. A “0” indicates that no class has been assigned to a particular dam, or that no dam exists.

The next section contains some watershed characteristics including the watershed area in acres and hectares and the land use composition of the watershed. A watershed is the entire area that will drain to a particular lake and is constrained by the topology and hydrology of the land. The watershed area was calculated by the US Geological Survey “StreamStats” program. This area map was then used to calculate land uses from the most recent (2011) National Land Use Cover data on the NYSDEC ArcGIS mapping program. The map itself is shown on the left side of the front page. In general, blue colors show water, green and light brown show forested or shrub land, yellow and dark brown are agriculture, and pink to red is developed land. The program participation section lists the years the lake has been sampled through CSLAP and the names of the 2016 samplers.

The next section includes four boxes. The trophic state of a lake refers to its nutrient loading and productivity- in other words, how much algae is produced, and the cause (nutrients) and outcome (changes in clarity) of this algae growth. An oligotrophic lake has low nutrient and algae levels (low productivity) and high clarity while a eutrophic lake has high nutrient and algae levels (high productivity) and low clarity. Mesotrophic lakes fall somewhere in the middle. For most lakes, the nutrient of concern is phosphorus. A more productive lake will

support more plant life, which may be good for warmwater fish, but may lower the quality of the lake if growth becomes excessive.

The harmful algal bloom susceptibility section contains a summary of the available historical HAB data. Although the factors that lead to the formation of HAB's is not yet well-understood, a history of HAB occurrences and high nutrient levels may indicate a susceptibility in the lake that could result in more HAB events in the future.

The invasive vulnerability section indicates if aquatic invasive species (AIS) are found in this lake or in nearby lakes. Invasive species are non-native and tend to rapidly colonize a waterbody once introduced, leaving little space for native species. Lakes with invasives or near other lakes with invasives are vulnerable to introductions of new AIS.

The next section is the priority waterbody list (PWL) assessment section. The PWL is a statewide inventory of the waters of New York State that DEC uses to track support (or impairment) of water uses, overall assessment water quality, causes and sources of water quality impact/impairment, and the status of restoration, protection and other water quality activities and efforts. A PWL assessment is broken into categories that include the following: potable water, swimming or public bathing, recreation, aquatic life, aesthetics, habitat, and fish consumption. All of the categories except aesthetics and habitat are assessed on a scale to determine if each of the listed uses are supported. The scale goes from best to worst in the following progression: fully supported, threatened, stressed, impaired, and precluded. Aesthetics and habitat are evaluated as good, fair, or poor. The cited PWL assessment reflects the "worst" assessment for the lake. The full PWL assessment for each lake can be found on the DEC website by searching on "PWL" and the lake basin, at <http://www.dec.ny.gov/chemical/36730.html#WIPWL>.

The rest of the report contains a collection of tables and charts. A glossary of all the water quality and HABs indicators used in the plots and tables is included below. Of particular note are the seasonal change and trend columns in the table. The long term trend column tells you if there is an increasing or decreasing trend, or no change, over time- one arrow equals a weak trend, and two arrows equals a strong trend. This may not agree with the seasonal change sparkline chart, which only shows the 2016 summer trends. Whether an increasing or decreasing trend is good for the lake depends on the indicator being evaluated, but in general green is good, red is bad. .

The next table contains a summary of open water and shoreline HABs data for the lake, along with the associated HAB notification information. Open water (mid lake) samples are collected routinely during each CSLAP sampling session. If a HAB is suspected, a sample from the worst part of the bloom (usually along the shoreline) is collected and sent in for laboratory confirmation. A HAB notification is added to the HAB database where entries are updated on a weekly basis. Additional information- samples or visual reports- are used to update these listings. The data graphs include the World Health Organization (WHO) high risk criteria to protect swimmers and the EPA low risk criteria to protect those using *treated* (not raw lake) water for drinking.

The Long Term Trend Analysis includes graphs showing the summer (mid-June thru mid-September) average for several of the key CSLAP water quality indicators, for each of the

years the lake was sampled through CSLAP. The graphs include relevant criteria (trophic categories, water quality standards,...) and boundaries separating these criteria.

The In-Season Analysis picks out two indicators- water temperature and water clarity- that are most frequently considered by lake residents as indicative of seasonal changes. These graphs are generated at any time during the sampling season for any CSLAP samplers that enters CSLAP data into the NYSFOLA on-line data entry program. The plots in this report show 2016 data compared to the normal seasonal variability for this lake.

The next section of the report includes Lake Use Scorecard. The scorecard presents the results of the existing Priority Waterbody List assessment for this lake in a graphical form and compares it to information from the current year and average values from CSLAP data and other lake information. The scorecard also includes a column that lists some primary issues that could impact specific use categories. Multiple issues could affect each designated use, but only the primary issue is listed.

The final section of the report is the Lake Summary. This includes a brief summary of the 2016 and historical CSLAP data for the lake. It is essentially the same as the Q&A section of the previous CSLAP reports, and with the Lake Use Scorecard, represents perhaps the most easily understood single page summary of the CSLAP data for the lake. This was intentionally created as the last page of the report to allow easy copying and distribution to lake association members, neighbors and others interested in the condition of the lake and the results from the CSLAP sampling.

Glossary of water quality and HAB indicators

Clarity (m): The depth to which a Secchi disk lowered into the water is visible, measured in meters. Water clarity is one of the trophic indicators for each lake.

TP (mg/l): Total phosphorus, measured in milligrams per liter at the lake surface (1.5 meters below the surface). TP includes all dissolved and particulate forms of phosphorus.

Deep TP: Total phosphorus measured in milligrams per liter at depth (1-2 meters above the lake bottom at the deepest part of the lake)

TN: Total nitrogen, measured in milligrams per liter at the lake surface. TN includes all forms of nitrogen, including **NO_x** (nitrite and nitrate) and **NH₄** (ammonia).

N:P Ratio: The ratio of total nitrogen to total phosphorus, unitless (mass ratio). This ratio helps determine if a lake is phosphorous or nitrogen limited.

Chl.a (ug/l): Chlorophyll a, measured in micrograms per liter

pH: A range from 0 to 14, with 0 being the most acidic and 14 being the most basic or alkaline. A healthy lake generally ranges between 6 and 9.

Cond (umho/cm): Specific conductance is a measure of the conductivity of water. A higher value indicates the presence of more dissolved ions that help conduct electricity. Conductivity results may indicate hard or softwater conditions with high ion concentrations resulting in hardwater.

Upper Temp (degC): Surface temperature, measured in degrees Celsius

Deep Temp (degC): Bottom temperature, measured in degrees Celsius

BG Chl.a (ug/L): Chlorophyll a from blue-green algae, measured in micrograms per liter

HABs Reported?: Were any algal blooms reported within a week of the dates listed, and, if so, were they located along the shoreline, in open water, or both?

BGA: Blue-green algae

Microcystin: The most common HAB liver toxin; total microcystin above 20 micrograms per liter indicates a “high toxin” bloom. However, ALL BGA blooms should be avoided, even if toxin levels are low.

Anatoxin-a: Another type of toxin that may be produced in a HAB and may be more dangerous as it targets the central nervous system. Neither EPA nor NYS has developed a risk threshold for anatoxin-a, although readings above 4 micrograms per liter are believed to represent an elevated risk.