



HAB Status


2017 Open Water Algae Samples



2017 Shoreline Algae Samples



2017 Open Water Toxin Levels




2017 Shoreline Toxin Levels




Canadarago Lake Long Term Trend Analysis


Clarity




Chlorophyll a



Surface and Deep Phosphorus




Lake Perception




Canadarago Lake Long Term Trend Analysis


Nitrogen




pH



Temperature




Specific Conductance




Canadarago Lake In-Season Analysis

In Season Temperature
























In Season Water Clarity



Scorecard

Lake Use

| | PWL | Average Year | 2017 | Primary Issue |
|------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|-----------------|
| Potable Water |  |  |  | Algae levels |
| Swimming |  |  |  | Algae blooms |
| Recreation |  |  |  | Algae levels |
| Aquatic Life |  |  |  | Bottom oxygen |
| Aesthetics |  |  |  | Poor perception |
| Habitat |  |  |  | Invasive plants |
| Fish Consumption |  |  |  | Not applicable |

-  Supported/Good
-  Threatened/Fair
-  Stressed/Poor
-  Impaired
-  Not Known

Summary

2017 compared to prior years: Canadarago Lake was sampled through CSLAP from 1989 to 1993 and in 2017. The lake continues to be *mesotrophic*, or moderately unproductive, based on intermediate levels water clarity, nutrients (phosphorus) and algae (chlorophyll *a*) levels. Water transparency was slightly lower than usual in 2017, likely due to slightly higher chlorophyll *a* readings, which in turn was consistent with slightly higher than normal surface and bottom phosphorus readings. However, despite the slightly lower water clarity, water quality perception was about as favorable as usual, although there was more extensive surface coverage of aquatic plants. Conductivity was lower than usual, consistent with slightly lower pH. Surface temperatures were slightly higher than usual in 2017.

Compared to nearby lakes: Canadarago Lake has similar water clarity, nutrient and algae levels, compared to other nearby (Central region) lakes, although it is slightly more productive than other nearby larger lakes. Aquatic plant coverage is usually slightly lower than in many of these other lakes, but was more extensive in 2017. It is not known if this is due to surface growth of invasive or native plants. Chloride levels are between the 25th and 50th percentile for New York state lakes, indicating a potential for aquatic life impacts from road salt (although no impacts have been reported or documented).

Trends: Phosphorus readings were significantly higher than in the period from 1989 to 1993. pH readings may have increased slightly over the last 25 years. Plant coverage may be increasing, and conductivity may be decreasing, but these changes are not (yet) statistically significant.

Algal blooms and HABS: Water quality conditions indicated a low to moderate susceptibility to blooms, although some *Dolichospermum* (previously called *Anabaena*) shoreline blooms were reported in 2015 and 2016. The algae community in the open water samples is comprised of several different taxa, particularly *Cryptophytes*, and overall algae levels are low. Open water and shoreline bloom toxins have not been detectable in any samples. No shoreline or open water blooms were apparent in 2017.

Aquatic invasive species: Eurasian watermilfoil, curly leafed pondweed and starry stonewort are found in Canadarago Lake, indicating an overall high vulnerability to AIS. This may be consistent with ready public access to the lake. Calcium levels are high enough to support zebra mussels, and these invasive mussels have been found in Canadarago Lake (along with rusty crayfish and common carp).

Indicated Actions: Individual stewardship activities such as pumping your septic system, growing a buffer of native plants next to the water bodies, and reducing erosion from shoreline properties and runoff into the lake will help to improve lake health by reducing nutrient and sediment loading to the lake. Visiting boats should be inspected to reduce the risk of new invasive species, and continued monitoring for invasive species is warranted. Continued algae bloom education and monitoring for HABS is recommended, particularly since shoreline blooms are periodically reported in Canadarago Lake.

How to Read the Report

This guide provides a description of the CSLAP report by section and a glossary. The sampling site is indicated in the header for lakes with more than one routine sampling site.

Physical Characteristics influence lake quality:

- Surface area is the lake's surface in acres and hectares.
- Max depth is the water depth measured at the deepest part of the lake in feet and meters.
- Mean depth is either known from lake bathymetry or is 0.46 of the maximum depth.
- Retention time is the time it takes for water to pass through a lake in years. This indicates the influence of the watershed on lake conditions.
- Lake classification describes the “best uses” for this lake. Class AA, AAspec, and A lakes may be used as sources of potable water. Class B lakes are suitable for contact recreational activities, like swimming. Class C lakes are suitable for non-contact recreational activities, including fishing, although they may still support swimming. The addition of a T or TS to any of these classes indicates the ability of a lake to support trout populations and/or trout spawning.
- Dam classification defines the hazard class of a dam. Class A, B, C, and D dams are defined as low, intermediate, high, or negligible/no hazard dams in that order. “0” indicates that no class has been assigned to a particular dam, or that no dam exists.

Watershed characteristics influence lake water quality:

- Watershed area in acres and hectares
- Land use data come from the most recent (2011) US Geological Survey National Land Use Cover dataset

CSLAP Participation lists the sampling years and the current year volunteers.

Key lake status indicators summarize lake conditions:

- Trophic state of a lake refers to its nutrient loading and productivity, measured by phosphorus, algae, and clarity. An oligotrophic lake has low nutrient and algae levels (low productivity) and high clarity while a eutrophic lake has high nutrient and algae levels (high productivity) and low clarity. Mesotrophic lakes fall in the middle.
- Harmful algal bloom susceptibility summarizes the available historical HAB data and indicates the potential for future HAB events.
- Invasive vulnerability indicates whether aquatic invasive species are found in this lake or in nearby lakes, indicating the potential for further introductions.
- Priority waterbody list (PWL) assessment is based on the assessment of use categories and summarized as fully supported, threatened, stressed,

impaired, or precluded. Aesthetics and habitat are evaluated as good, fair, or poor. The cited PWL assessment reflects the “worst” assessment for the lake. The full PWL assessment can be found at <http://www.dec.ny.gov/chemical/36730.html#WIPWL>.

Current year sampling results

- Results for each of the sampling sessions in the year are in tabular form. The seasonal change graphically shows the current year results. Red shading indicates eutrophic readings.
- HAB notification periods on the DEC website, updated weekly <http://www.dec.ny.gov/chemical/83310.html>
- Shoreline HAB sample dates and results. Samples are collected from the area that appears to have the worst bloom. Red shading indicates a confirmed HAB.
- HAB sample algae analysis. Algae types typically change during the season. These charts show the amount of the different types of algae found in each mid-lake or shoreline sample. Samples with high levels of BGA are HABs. The second set of charts show the level of toxins found in open water and shoreline samples compared to the World Health Organization (WHO) guidelines.
- If there are more than ten shoreline bloom samples collected in a year, bloom sample information is instead summarized by month (May-Oct.) as minimum, average, and maximum values for blue-green algae and microcystin.

Long Term Trend Analysis puts the current year findings in context. Summer averages (mid-June thru mid-September) for each of the CSLAP years show trends in key water quality indicators. The graphs include relevant criteria (trophic categories, water quality standards, etc.) and boundaries separating these criteria.

In-Season Analysis shows water temperature and water clarity during the sampling season. These indicate seasonal changes and show the sample year results compared to the typical historical readings for those dates.

The Lake Use Scorecard presents the results of the existing Priority Waterbody List assessment for this lake in a graphical form and compares it to information from the current year and average values from CSLAP data and other lake information. Primary issues that could impact specific use categories are identified, although more issues could also affect each designated use.

The Lake Summary reviews and encapsulates the data in the lake report, and provides suggested actions for lake management.

Glossary of water quality and HAB indicators

Clarity (m): The depth to which a Secchi disk lowered into the water is visible, measured in meters. Water clarity is one of the trophic indicators for each lake.

TP (mg/L): Total phosphorus, measured in milligrams per liter at the lake surface (1.5 meters below the surface). TP includes all dissolved and particulate forms of phosphorus.

Deep TP: Total phosphorus measured in milligrams per liter at depth (1-2 meters above the lake bottom at the deepest part of the lake)

TN: Total nitrogen, measured in milligrams per liter at the lake surface. TN includes all forms of nitrogen, including **NO_x** (nitrite and nitrate) and **NH₄** (ammonia).

N:P Ratio: The ratio of total nitrogen to total phosphorus, unitless (mass ratio). This ratio helps determine if a lake is phosphorous or nitrogen limited.

Chl.a (µg/L): Chlorophyll a, measured in micrograms per liter. Indicates the amount of algae in the water column.

pH: A range from 0 to 14, with 0 being the most acidic and 14 being the most basic or alkaline. A healthy lake generally ranges between 6.5 and 8.5.

Cond (µmho/cm): Specific conductance is a measure of the conductivity of water. A higher value indicates the presence of more dissolved ions. High ion concentrations indicate hardwater, and low show softwater.

Upper Temp (°C): Surface temperature, measured in degrees Celsius

Deep Temp (°C): Bottom temperature, measured in degrees Celsius

BG Chl.a (µg/L): Chlorophyll a from blue-green algae, measured in micrograms per liter

HABs: Harmful Algal Blooms. Algal blooms that have the appearance of cyanobacteria (BGA)

BGA: Blue-green algae, also known as cyanobacteria

Microcystin (µg/L): The most common HAB liver toxin; total microcystin above 20 micrograms per liter indicates a “high toxin” bloom. However, ALL BGA blooms should be avoided, even if toxin levels are low.

Anatoxin-a (µg/L): A toxin that may be produced in a HAB which targets the central nervous system. Neither EPA nor NYS has developed a risk threshold for anatoxin-a, although readings above 4 micrograms per liter are believed to represent an elevated risk.