












<b>Lake Characteristics</b>	Surface Area (ac/ha)	454	184
	Max Depth (ft/m)	49	15
	Mean Depth (ft/m)	17	5
	Retention Time (years)	0.33	
	Water Class	B	
	Dam Class	C	
<b>Watershed Characteristics</b>	Watershed Area(ac/ha)	16244	6574
	Watershed/Lake Ratio	36	
	Lake and Wetlands	3.4%	
	Agricultural	22.6%	
	Forests, shrubs, grasses	71.6%	
	Residential	2.4%	
	Urban	0.0%	
<b>CSLAP Participation</b>	Years	1986-1988, 1990-1991, 1998-2017	
	Volunteers	Scott Barrey	

Trophic State	HABs Susceptibility	Invasive Vulnerability	PWL Assessment
Mesoeutrophic	Moderate	High	Stressed


Open Water Indicators	2017 Sampling Results								Seasonal Change	Long Term Avg.
	6/13	7/11	7/25	8/8	8/15	8/22	9/6	9/19		
Chl.a (µg/L)	6.8	4.5	8.1	9.9	11.5	8.7	18.7	8.9		12.0
BG Chl.a (µg/L)	2.0	3.2	3.7	0.5	7.8	6.3	7.0	4.0		3.5
Clarity (m)	3.8	3	2.9	3	2.4	2.2	2.2	3.3		3.0
pH	6.8	8.2	7.9	7.2	7.7	8.3	7.9	8.1		7.7
Cond (µmho/cm)	114.6	91.2	108.2	115.4	104.3	120.3	105.4	111.7		120
Surf Temp (°C)	25	25	25	25	25	25	22	25		22
Bott Temp (°C)	19	16	16	16	15	16	14	14		14
TN (mg/L)	.507	.261	.528	.255	.247	.33	.289	.295		0.492
TP (mg/L)	.016	.017	.017	.021	.019	.023	.017	.018		0.019
Deep TP (mg/L)	.044	.072	.139	.433	.621	.489	1.929	.86		0.268
N:P Ratio	32	15	31	12	13	14	17	16		

Date of first listing	Date of last listing	# of weeks on DEC notification list	# of weeks with updates
		0	


Shoreline HAB Sample Dates 2017											
HAB Indicators	HAB Criteria										
BGA	25 µg/L	NA									
Microcystin	20 µg/L	NA									
Anatoxin-a		NA									

## HAB Status


### 2017 Open Water Algae Samples



### 2017 Shoreline Algae Samples



### 2017 Open Water Toxin Levels




### 2017 Shoreline Toxin Levels




## Cuba Lake Long Term Trend Analysis


### Clarity




### Chlorophyll a



### Surface and Deep Phosphorus




### Lake Perception




# Cuba Lake Long Term Trend Analysis


## Nitrogen




## pH



## Temperature




## Specific Conductance




# Cuba Lake In-Season Analysis

## In Season Temperature
























## In Season Water Clarity



# Scorecard

## Lake Use

Potable Water				No impacts
Swimming				Algae blooms
Recreation				Algae levels
Aquatic Life				Bottom Oxygen
Aesthetics				Algae blooms
Habitat				Invasive plants
Fish Consumption				Not applicable

PWL

Average  
Year

2017

Primary Issue



## Summary

**2017 compared to prior years:** Cuba Lake continues to be typical of *mesoeutrophic* (moderately productive) lakes, based on high algae levels and intermediate nutrient and water clarity readings. Conductivity and total nitrogen levels were lower than usual in 2017, while water temperatures were higher.

**Compared to nearby lakes:** Cuba Lake has slightly higher water clarity and lower nutrient and algae levels than most Western region lakes. Chloride levels are between the 25<sup>th</sup> and 50<sup>th</sup> percentile levels for New York state lakes, indicating little potential for aquatic life impacts from road salt.

**Trends:** No clear long-term trends have been apparent in the CSLAP data. Surface and bottom phosphorus readings have increased slightly, while pH has decreased slightly. However, none of these changes are statistically significant.

**Algal blooms and HABS:** Water quality conditions indicated a moderate susceptibility to blooms, and shoreline water blooms are periodically reported. Blue green algae levels do rise slightly during the summer, but both open water and shoreline bloom samples have toxin levels that are typically not detectable. No blooms were reported in 2017.

**Aquatic invasive species:** Eurasian watermilfoil, curly-leafed pondweed, brittle naiad, and common carp have been documented in Cuba Lake, and the public access and proximity to other nearby lakes with AIS indicate a high vulnerability to new AIS introductions.

**Indicated Actions:** Individual stewardship activities such as pumping your septic system, growing a buffer of native plants next to the water bodies, and reducing erosion from shoreline properties and runoff into the lake will help to improve lake health by reducing nutrient and sediment loading to the lake. Visiting boats should be inspected to reduce the risk of new invasive species, and continued monitoring for invasive species is warranted. Continued algae bloom education and monitoring for HABS is recommended, since many western New York lakes are susceptible to these blooms.

## How to Read the Report

This guide provides a description of the CSLAP report by section and a glossary. The sampling site is indicated in the header for lakes with more than one routine sampling site.

### **Physical Characteristics influence lake quality:**

- Surface area is the lake's surface in acres and hectares.
- Max depth is the water depth measured at the deepest part of the lake in feet and meters.
- Mean depth is either known from lake bathymetry or is 0.46 of the maximum depth.
- Retention time is the time it takes for water to pass through a lake in years. This indicates the influence of the watershed on lake conditions.
- Lake classification describes the “best uses” for this lake. Class AA, AAspec, and A lakes may be used as sources of potable water. Class B lakes are suitable for contact recreational activities, like swimming. Class C lakes are suitable for non-contact recreational activities, including fishing, although they may still support swimming. The addition of a T or TS to any of these classes indicates the ability of a lake to support trout populations and/or trout spawning.
- Dam classification defines the hazard class of a dam. Class A, B, C, and D dams are defined as low, intermediate, high, or negligible/no hazard dams in that order. “0” indicates that no class has been assigned to a particular dam, or that no dam exists.

### **Watershed characteristics influence lake water quality:**

- Watershed area in acres and hectares
- Land use data come from the most recent (2011) US Geological Survey National Land Use Cover dataset

**CSLAP Participation** lists the sampling years and the current year volunteers.

### **Key lake status indicators summarize lake conditions:**

- Trophic state of a lake refers to its nutrient loading and productivity, measured by phosphorus, algae, and clarity. An oligotrophic lake has low nutrient and algae levels (low productivity) and high clarity while a eutrophic lake has high nutrient and algae levels (high productivity) and low clarity. Mesotrophic lakes fall in the middle.
- Harmful algal bloom susceptibility summarizes the available historical HAB data and indicates the potential for future HAB events.
- Invasive vulnerability indicates whether aquatic invasive species are found in this lake or in nearby lakes, indicating the potential for further introductions.
- Priority waterbody list (PWL) assessment is based on the assessment of use categories and summarized as fully supported, threatened, stressed,

impaired, or precluded. Aesthetics and habitat are evaluated as good, fair, or poor. The cited PWL assessment reflects the “worst” assessment for the lake. The full PWL assessment can be found at <http://www.dec.ny.gov/chemical/36730.html#WIPWL>.

### **Current year sampling results**

- Results for each of the sampling sessions in the year are in tabular form. The seasonal change graphically shows the current year results. Red shading indicates eutrophic readings.
- HAB notification periods on the DEC website, updated weekly <http://www.dec.ny.gov/chemical/83310.html>
- Shoreline HAB sample dates and results. Samples are collected from the area that appears to have the worst bloom. Red shading indicates a confirmed HAB.
- HAB sample algae analysis. Algae types typically change during the season. These charts show the amount of the different types of algae found in each mid-lake or shoreline sample. Samples with high levels of BGA are HABs. The second set of charts show the level of toxins found in open water and shoreline samples compared to the World Health Organization (WHO) guidelines.
- If there are more than ten shoreline bloom samples collected in a year, bloom sample information is instead summarized by month (May-Oct.) as minimum, average, and maximum values for blue-green algae and microcystin.

**Long Term Trend Analysis** puts the current year findings in context. Summer averages (mid-June thru mid-September) for each of the CSLAP years show trends in key water quality indicators. The graphs include relevant criteria (trophic categories, water quality standards, etc.) and boundaries separating these criteria.

**In-Season Analysis** shows water temperature and water clarity during the sampling season. These indicate seasonal changes and show the sample year results compared to the typical historical readings for those dates.

**The Lake Use Scorecard** presents the results of the existing Priority Waterbody List assessment for this lake in a graphical form and compares it to information from the current year and average values from CSLAP data and other lake information. Primary issues that could impact specific use categories are identified, although more issues could also affect each designated use.

**The Lake Summary** reviews and encapsulates the data in the lake report, and provides suggested actions for lake management.

## Glossary of water quality and HAB indicators

**Clarity (m):** The depth to which a Secchi disk lowered into the water is visible, measured in meters. Water clarity is one of the trophic indicators for each lake.

**TP (mg/L):** Total phosphorus, measured in milligrams per liter at the lake surface (1.5 meters below the surface). TP includes all dissolved and particulate forms of phosphorus.

**Deep TP:** Total phosphorus measured in milligrams per liter at depth (1-2 meters above the lake bottom at the deepest part of the lake)

**TN:** Total nitrogen, measured in milligrams per liter at the lake surface. TN includes all forms of nitrogen, including **NO<sub>x</sub>** (nitrite and nitrate) and **NH<sub>4</sub>** (ammonia).

**N:P Ratio:** The ratio of total nitrogen to total phosphorus, unitless (mass ratio). This ratio helps determine if a lake is phosphorous or nitrogen limited.

**Chl.a (µg/L):** Chlorophyll a, measured in micrograms per liter. Indicates the amount of algae in the water column.

**pH:** A range from 0 to 14, with 0 being the most acidic and 14 being the most basic or alkaline. A healthy lake generally ranges between 6.5 and 8.5.

**Cond (µmho/cm):** Specific conductance is a measure of the conductivity of water. A higher value indicates the presence of more dissolved ions. High ion concentrations indicate hardwater, and low show softwater.

**Upper Temp (°C):** Surface temperature, measured in degrees Celsius

**Deep Temp (°C):** Bottom temperature, measured in degrees Celsius

**BG Chl.a (µg/L):** Chlorophyll a from blue-green algae, measured in micrograms per liter

**HABs: Harmful Algal Blooms.** Algal blooms that have the appearance of cyanobacteria (BGA)

**BGA:** Blue-green algae, also known as cyanobacteria

**Microcystin (µg/L):** The most common HAB liver toxin; total microcystin above 20 micrograms per liter indicates a “high toxin” bloom. However, ALL BGA blooms should be avoided, even if toxin levels are low.

**Anatoxin-a (µg/L):** A toxin that may be produced in a HAB which targets the central nervous system. Neither EPA nor NYS has developed a risk threshold for anatoxin-a, although readings above 4 micrograms per liter are believed to represent an elevated risk.