

Species Status Assessment

Common Name: a giant casemaker caddisfly

Date:

Scientific Name: *Banksiola calva*

Proposed By: Luke Myers

Class: Insecta

Family: Phyragenidae

Species Synopsis (a short paragraph which describes species taxonomy, distribution, recent trends, and habitat in New York):

This species is known to inhabit bogs and has been identified in St. Lawrence and Franklin counties. (Myers et al. 2010). Outside of New York, this species has been collected in Massachusetts, New Hampshire, and Connecticut (Banks, 1943; Morse and Blickle, 1953; Wiggins, 1998; Morse 2010).

I. Status

a. Current legal protected Status

i. **Federal:** Not listed _____ **Candidate:** _____

ii. **New York:** Not listed _____

b. Natural Heritage Program

i. **Global:** G2G3 _____

ii. **New York:** S1S2 _____ **Tracked by NYNHP?:** Yes _____

Other Ranks:

- New York 2025 SGCN status: Species of Greatest Conservation Need
- COSEWIC: Not listed
- IUCN Red List: Not listed
- Northeast Regional SGCN: Proposed RSGCN

Status Discussion:

II. Abundance and Distribution Trends

Region	Present?	Abundance	Distribution	Time Frame	Listing status	SGCN?
North America	Yes	Unknown	Unknown			-
Northeastern US	Yes	Unknown	Unknown			-
New York	Yes	Unknown	Unknown			Yes
Connecticut	No	-	-			-

Region	Present?	Abundance	Distribution	Time Frame	Listing status	SGCN?
Massachusetts	Yes	Unknown	Unknown			-
New Jersey	No	-	-			-
Pennsylvania	No	-	-			-
Vermont	No	-	-			-
Ontario	No	-	-			-
Quebec	No	-	-			-

Column options

Present?: Yes; No; Unknown; No data; (blank) or Choose an Item

Abundance and Distribution: Declining; Increasing; Stable; Unknown; Extirpated; N/A; (blank) or Choose an item

SGCN?: Yes; No; Unknown; (blank) or Choose an item

Monitoring in New York (specify any monitoring activities or regular surveys that are conducted in New York):

None

Trends Discussion (insert map of North American/regional distribution and status):




Figure 1. Conservation status of *Banksiola calva* in North America (NatureServe 2024.)

III. New York Rarity (provide map, numbers, and percent of state occupied)




Figure 2. Distribution of *Banksiola calva* in New York State (Myers et al. 2010).

Figure 4+. (Any additional maps, charts or tables (title, source, date))

Details of historic and current occurrence:

New York’s Contribution to Species North American Range:

Percent of North American Range in NY	Classification of NY Range	Distance to core population, if not in NY
Choose an item.	Choose an item.	

Column options

Percent of North American Range in NY: 100% (endemic); 76-99%; 51-75%; 26-50%; 1-25%; 0%; Choose an item

Classification of NY Range: Core; Peripheral; Disjunct; (blank) or Choose an item

IV. Primary Habitat or Community Type (from NY crosswalk of NE Aquatic, Marine, or Terrestrial Habitat Classification Systems):

- a. Lacustrine: shallow water
- b. Palustrine: Bog/fen, herbaceous wetland

c. Riverine: low gradient, pool

Habitat or Community Type Trend in New York

Habitat Specialist?	Indicator Species?	Habitat/Community Trend	Time frame of Decline/Increase
Choose an item.	Choose an item.	Choose an item.	

Column options

Habitat Specialist and Indicator Species: Yes; No; Unknown; (blank) or Choose an item

Habitat/Community Trend: Declining; Stable; Increasing; Unknown; (blank) or Choose an item

Habitat Discussion:

V. Species Demographic, and Life History:

Breeder in NY?	Non-breeder in NY?	Migratory Only?	Summer Resident?	Winter Resident?	Anadromous/Catadromous?
-	-	-	-	-	-

Column options

First 5 fields: Yes; No; Unknown; (blank) or Choose an item

Anadromous/Catadromous: Anadromous; Catadromous; (blank) or Choose an item

Species Demographics and Life History Discussion *(include information about species life span, reproductive longevity, reproductive capacity, age to maturity, and ability to disperse and colonize):*

VI. Threats (from NY 2015 SWAP or newly described):

Threat Level 1	Threat Level 2	Threat Level 3	Spatial Extent	Severity	Immediacy	Trend	Certainty
4. Transportation & Service Corridors	4.1 Roads & Railroads	4.1.1 Roads (runoff)	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
4. Transportation & Service Corridors	4.1 Roads & Railroads	4.1.3 Bridges (construction/maintenance)	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
5. Biological Resource Use	5.3 Logging & Wood Harvesting	-	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
7. Natural System Modifications	7.2 Dams & Water Management/Use	7.2.1 Water level management using dams (altered hydrology, sedimentation)	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
8. Invasive & Other Problematic Species	8.1 Invasive Non-Native Plants & Animals	-	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
9. Pollution	9.1 Domestic & Urban Wastewater	-	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
9. Pollution	9.2 Industrial & Military Effluents	-	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
9. Pollution	9.3 Agricultural & Forestry Effluents	9.3.1 Nutrient loads (run off, algal blooms)	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
11. Climate Change	11.1 Habitat Shifting & Alteration	-	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.
11. Climate Change	11.4 Changes in Precipitation & Hydrological Regimes	11.4.2 Droughts	Choose an item.	Choose an item.	Choose an item.	Choose an item.	Choose an item.

Table 2. Threats to *Banksiola calva*.

Any activity which might lead to water contamination, siltation, warming of waterways, or the alteration of natural hydrology could directly and indirectly impact riparian habitats and *Banksiola calva* populations. Such threats might include roadway and agricultural runoff, industrial pollution, dams, bridge construction and maintenance, logging activities, and development near riparian habitats (NYS DEC 2005). In addition, siltation decreases the amount of sunlight that reaches aquatic plants (EPA 2005) and lowers the quality of habitats needed for a variety of aquatic species (NYS DEC 2005). Point source pollution, such as effluents from municipal and industrial facilities, contribute to the degradation and pollution of aquatic habitats (EPA 2022, NYS DEC 2005, Mahar and Landry 2013, Strayer et al. 2004).

While modern day agricultural and silvicultural practices are an important aspect of the New York State economy, it is important to consider the effects on ecosystems and species. As these practices move closer to rivers, the natural riparian buffers are often removed. Furthermore, they slow or reduce runoff from farm fields and pastures, such as contaminants from pesticides, fertilizers, manure, and sludge, into waterways (EPA 2005, NYS DEC 2005, Souza et al. 2020). Excessive fertilizer use can lead to algal blooms that can be deadly to aquatic life and overgrazing of livestock in fields could introduce pathogens, oxygen-demanding organics and solids, and invasive species to aquatic ecosystems (EPA 2005).

Approximately 10% of introduced, non-native species could have an impact on the health of ecosystems (McCormick et al. 2009). Invasive plants tend to outcompete native plants and can change natural processes (NYS DEC 2005). There is an increased risk of runoff and erosion when these plants are along streams and rivers. Aquatic invasive plants and animals can alter the water chemistry, change the nutrient regime, or decrease the dissolved oxygen levels. Introduced fish can alter trophic relationships resulting in a change in native fish populations and decreased water quality (McCormick et al. 2009).

Climate change is another threat that is likely to have lasting effects on riverine systems. Irregular weather patterns can cause extreme drought, flooding, and temperature fluctuations. Heat waves are expected to be more intense (Frankson et al. 2022). The Northeast Region of the United States is expected to experience an increase in precipitation, more frequent storms, and higher than normal temperatures (EPA 2016, EPA 2022). Precipitation is expected to increase 10% to 15% in southern New York and 15 to 20% in northern New York by 2050 (Frankson et al. 2022). Extreme flooding can cause widespread erosion and runoff with added risk of contamination if flooding occurs at remediation sites, industrial sites, or wastewater treatment facilities (EPA 2016, EPA 2022). Temperature increases can significantly alter ecosystems. As water temperatures rise, the amount of dissolved oxygen decreases and evaporation increases, potentially lowering lake and stream levels (EPA 2022). Any combination of these events could change species distributions (EPA 2022) and those that cannot adapt or migrate may be extirpated from some areas (NYS DEC 2005).

Are there regulatory mechanisms that protect the species or its habitat in New York?

Yes:

No:

Unknown:

If yes, describe mechanism and whether adequate to protect species/habitat:

The Protection of Waters Program provides protection for rivers, streams, lakes, and ponds under Article 15 of the NYS Environmental Conservation Law.

Describe knowledge of management/conservation actions that are needed for recovery/conservation, or to eliminate, minimize, or compensate for the identified threats:

Protect water quality and reduce contamination and hydrological alteration (such as agricultural or road runoff, shoreline development, and damming) (NYS DEC 2005). Protect stream quality by maintaining both a riparian buffer that includes herbaceous and/or woody vegetation along the shoreline, and a significant forested buffer. These buffers reduce sediment and contaminant runoff (EPA 2005, NYS DEC 2005, Souza et al. 2020), provide shade, regulate temperature, and provide organic matter to animals (Hughes and Vadas 2021). Riparian zones with herbaceous and woody vegetation showed improved “indicator scores” for macroinvertebrates and fishes in the Midwest (Hughes and Vadas 2021).

Remove barriers to maintain or restore natural flow to waterways. Where removal is not possible, research alternatives that allow flow above and below a barrier.

In general, avoid stream crossings. If crossings are unavoidable, use Best Management Practices (BMP) to minimize disturbance to streams. Time periods of disturbance when water flow is low or normal and install stream-crossing structures at a right angle to the stream (Watershed Agricultural Council Forestry Program 2018). Temporary methods to reduce runoff include water bars, gravel, geotextile fabric, rubber belt deflectors, open top culverts, strawbales, silt fencing, control blankets, and straw wattles (Watershed Agricultural Council Forestry Program 2018). Restore the disturbed area with native species as soon as possible. Areas that have been logged may also need ruts to be smoothed to reduce surface runoff (Watershed Agricultural Council Forestry Program 2018). Hughes and Vadas (2021) suggest that Best Management Practices may need to be applied to entire stream lengths and catchments to fully restore an aquatic ecosystem. If this is not possible, restore or manage a larger area around the directly disturbed area.

In general, tailor agricultural management plans to local conditions (e.g., soils, slope, land use). Often these plans aim to reduce pollution and increase farm productivity, but incentives could also be used to encourage sustainable farming practices. Proper management typically reduces runoff by 20-90% (EPA 2005). Practices to reduce runoff and erosion include cover crops, crop rotation, tillage, mulching, terraces, diversion, and water and sediment control basins (NYS Soil and Water Conservation Committee 2023). Consider using Integrated Pest Management (IPM) as an alternative to pesticide use. If pesticides and fertilizers are used, they should only be applied when needed, in the proper amount, and timed appropriately. In addition, rotate livestock to avoid overgrazing and to allow for vegetation regrowth. In heavy use areas, consider a runoff management system that reduces contamination of nearby streams and soils (EPA 2005, NYS Soil and Water Conservation Committee 2023). If needed, provide alternative water sources and shade to keep animals out of sensitive areas (EPA 2005). Large livestock farms may benefit from a manure waste treatment system, such as, anaerobic digestion, liquid/solid separation, composting, biological and chemical amendments, and waste facility cover (NYS Soil and Water Conservation Committee 2023).

Invasive species management can be time consuming and costly. Reduce the likelihood of non-native species being introduced into waterways. Boat-washing stations at boat launches can reduce transport of invasive plants and animals to new waterbodies. Educate anglers about the risk of releasing unused baitfish. If vulnerable species are present, consider a baitfish ban. Mechanical removal of some invasive plants may be needed in some rivers and streams. The use of pesticides to remove invasives can have a negative effect on ecosystems (McCormick et al. 2009) and should be a last resort to control invasive species.

Climate change is a global challenge. However, there are local actions that can help mitigate extreme weather events. Industrial and municipal infrastructure should be improved or replaced to be more resilient to flooding events (EPA 2016, NYS Comptroller 2023). Some suggested actions include installing or improving pumps to remove floodwater from facilities and installing protective structures, such as floodwalls. Ensure that existing bridges, dams, levees, seawalls, retaining walls, and wind barriers are prepared for extreme weather (NYS Comptroller 2023). Decrease runoff and erosion severity by installing large culverts, planting vegetation along riverbanks, and protecting and restoring wetlands (EPA 2016, NYS Comptroller 2023).

Public outreach is an important tool for conservation. Many of the threats to aquatic habitats and species can have direct and indirect effects to human health and recreation. In addition, outreach and incentives for various industries would be valuable. Improvements to municipal and industrial treatment facilities was noted by NYS DEC (2005) and further improvements and plans to deal with spills should continue to be encouraged.

Action Category	Action	Description
A.1 Direct Habitat Management	A.1.1.1.2 Manual removal	Manually remove invasive plants
A.1 Direct Habitat Management	A.1.2.4.1 Plant for erosion management	Implement and maintain riparian buffers to reduce sedimentation and runoff
A.1 Direct Habitat Management	A.1.3.0.0 Mitigate human environmental impact	Implement temporary runoff reduction measures
A.1 Direct Habitat Management	A.1.3.1.0 Manage Access	Implement boat washing stations to reduce transport of invasives between waterbodies
A.1 Direct Habitat Management	A.1.3.3.0 Remove and improve anthropogenic infrastructure	Remove barriers to natural flow in waterways where feasible
A.1 Direct Habitat Management	A.1.3.3.0 Remove and improve anthropogenic infrastructure	Remove and improve industrial/municipal infrastructure to be more flood resilient
A.1 Direct Habitat Management	A.1.3.3.2 Culverts	Install culverts to reduce runoff and erosion severity
B.3 Outreach	B.3.1.0.0 Outreach, communication, and distribution	Educate anglers about the risks of releasing unused baitfish
B.5 Economic and Other Incentives	B.5.0.0.0 Economic and other incentives	Use incentives to encourage sustainable farming practices

Action Category	Action	Description
C.7 Legislative and Regulatory Framework or Tools	C.7.2.0.0 Create or amend policies, guidelines, or best practices	Consider baitfish bans in locations with vulnerable species
C.7 Legislative and Regulatory Framework or Tools	C.7.2.2.0 Create or amend best practices or guidelines	Use Best Management Practices when crossing streams, consider water flow when constructing in or around stream.

Table 3. Recommended conservation actions for *Banksiola calva*.

VII. References

Banks, N. 1943. Notes and Descriptions of Nearctic Trichoptera. Harvard University Museum of Comparative Zoology Bulletin 92: 341-369.

Blickle, R. L. and W. J. Morse, 1954. New species of Hydroptilidae (Trichoptera). Bulletin of the Brooklyn Entomological Society 49: 121-127.

Environmental Protection Agency (EPA). 2005. Protecting water quality from agricultural runoff. https://www.epa.gov/sites/default/files/2015-09/documents/ag_runoff_fact_sheet.pdf. Accessed on July 3, 2023.

Environmental Protection Agency (EPA). 2016. Adapting to climate change northeast. https://www.epa.gov/sites/default/files/2016-07/documents/northeast_fact_sheet.pdf. Accessed on August 10, 2023.

Environmental Protection Agency. 2022. Region 2 climate adaptation implementation plan. <https://www.epa.gov/system/files/documents/2022-10/bh508-R02%20Climate%20Adaptation%20Implementation%20Plan%209.19.2022%20HQ%20OP%20Copy.pdf>

Frankson, R., Kunkel, K.E., Champion, S.M., Stewart, B.C., Sweet, W, DeGaetano, A.T., & Spaccio, J. (2022). New York State Climate Summary 2022. National Oceanic and Atmospheric Administration National Centers for Environmental Information. <https://statesummaries.ncics.org/chapter/ny/>

Hughes, Robert M., and Robert L. Vadas Jr. 2021. Agricultural Effects on Streams and Rivers: A Western USA Focus. Water 13, no. 14: 1901. <https://doi.org/10.3390/w13141901>

Mahar, Amy and Jenny Landry. 2013. New York State Department of Environmental Conservation species status assessment for *Lasmigona subviridis* (Green Floater).

McCormick, Frank H., Glen C. Contreras, and Sherri L. Johnson. 2009. "Effects of nonindigenous invasive species on water quality and quantity." *A dynamic invasive species research vision: opportunities and priorities* 29 (2009): 111-120.

Morse, J. C. 2010. Trichoptera species of America north of Mexico. Available <http://entweb.clemson.edu/faculty/morseneartictrichopteraabbr.pdf> (Accessed, January 30, 2010).

Myers L.W., T.B. Mihuc and B.C. Kondratieff. 2010. Mayflies (Ephemeroptera), Stoneflies (Plecoptera), and Caddisflies (Trichoptera) of the Upper Hudson, Lake Champlain, and Northeastern Lake Ontario Watersheds: A baseline inventory with management considerations for SGCN and other rare and possibly imperiled species. Final Report to the New York State Department of Environmental Conservation.

NatureServe. 2024. NatureServe Explorer. Page last published 11/14/24. https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.119208/Banksiola_calva. Accessed November 21, 2024.

New York State Comptroller. 2023. New York's local governments adapting to climate change: challenges, solutions, and costs. <https://www.osc.state.ny.us/files/local-government/publications/pdf/climate-change-2023.pdf>

New York State Department of Environmental Conservation. 2005. A strategy for conserving New York's fish and wildlife resources. Final submission draft.

Souza, Francine N., Rodolfo Mariano, Tassio Moreira, and Sofia Campiolo. 2020. Influence of the landscape in different scales on the EPT community (Ephemeroptera, Plecoptera and Trichoptera) in the Atlantic Forest region. *Environmental monitoring and assessment* 129: 391-391.

Strayer, David L., J.A. Dowling, W.R. Haag, T.L. King, J.B. Layzer, T.J. Newton and S.J. Nichols. 2004. Changing perspectives on Pearly Mussels, North America's most Imperiled Animals. *BioScience* 54:429-439.

Watershed Agricultural Council Forestry Program. 2018. New York State forestry voluntary best management practices for water quality. <http://nysbmpguidelines.com/>. Accessed on June 20, 2023.

Wiggins, G. B. 1998. The caddisfly family Phryganeidae (Trichoptera). Royal Ontario Museum, Life Sciences Contributions. 306 pp.

Zaidel, Peter A., A. H. Roy, K. M. Houle, B. Lambert, B. H. Letcher, K. H. Nislow, C. Smith. 2021. Impacts of small dams on stream temperature. *Ecological indicators* 120:6-11.

Originally prepared by	
Date first prepared	
First revision	
Latest revision	11/21/2024 (Vincenzo Bonaiuto)