

# Species Status Assessment

**Common Name:** Mucket

**Date Updated:** 2/4/2024

**Scientific Name:** *Ortmanniana ligamentina* **Updated By:** A. Mahar & K. Crandall

**Class:** Bivalvia

**Family:** Unionidae

**Species Synopsis** (a short paragraph which describes species taxonomy, distribution, recent trends, and habitat in New York):

*Ortmanniana ligamentina* (syn. *Actinonaias ligamentina*, MolluscaBase eds., 2024) belongs to the subfamily *Ambleminae* and the tribe *Lampsilini*, which includes 17 extant and 6 likely extirpated New York species of the genera *Actinonaias*, *Epioblasma*, *Lampsilis*, *Leptodea*, *Ligumia*, *Obovaria*, *Potamilus*, *Ptychobranchnus*, *Toxolasma*, *Truncilla*, and *Villosa* (Haag 2012; Graf and Cummings 2011). *Ortmanniana ligamentina* is considered taxonomically synonymous with *Actinonaias ligamentina*. In 2021, the two North American *Actinonaias* species were reclassified into the *Ortmanniana* genus (Graf & Cummings, 2021). *O. Ligamentina* is named for its large, strong ligament (Watters et al. 2009).

This species typically inhabits fast flowing sections of large streams and rivers in cobble and gravel and is occasionally found in slow water and large lakes (Strayer & Jirka 1997; Williams et al., 2008). Since 1970, it has been found in ten New York waterbodies. It is often the most abundant mussel in the Allegheny River system's medium gradient streams (The Nature Conservancy, 2009). In addition to the upper Allegheny basin and Conewango Creek basin, *O. ligamentina* is also found in the French Creek and Lake Erie basins.

Although ranked as "imperiled" in New York at the edge of its range, this species is considered secure throughout its range. In North America, approximately 2/3 to ¾ of native mussel species are extinct, listed as endangered or threatened, or are in need of conservation status (Williams et al. 1993; Stein et al. 2000). While population trends in New York are unknown, based on sparse historical information, it is assumed that they too are declining due to a myriad of environmental stressors.

## I. Status

### a. Current legal protected Status

i. **Federal:** None **Candidate:** No

ii. **New York:** None;SGCN

### b. Natural Heritage Program

i. **Global:** G5 - Secure

ii. **New York:** S1S2 - Imperiled **Tracked by NYNHP?:** Yes

### Other Ranks:

-IUCN Red List: Least Concern (2011)

-Northeast Regional SGCN: No (2023)

-American Fisheries Society Status: Currently Stable (1993)

### Status Discussion:

This species is widely distributed and found throughout the Mississippi River system, with the exception of extreme southern and western reaches. It also occurs in the St. Lawrence River basin and tributaries of Lakes Erie, Michigan, and Ontario and is considered stable throughout much of its range and is globally secure (NatureServe 2013).

## II. Abundance and Distribution Trends

Region	Present?	Abundance	Distribution	Time Frame	Listing status	SGCN?
<b>North America</b>	Yes	Stable	Stable	Pre-2011		(blank)
<b>Northeastern US</b>	Yes	-	-			No
<b>New York</b>	Yes	-	-		S2	Yes
<b>Connecticut</b>	No	N/A	N/A			No
<b>Massachusetts</b>	No	N/A	N/A			No
<b>New Jersey</b>	No	N/A	N/A			No
<b>Pennsylvania</b>	Yes	-	-		S4	No
<b>Vermont</b>	No	N/A	N/A			No
<b>Ontario</b>	Yes	-	-		S4	(blank)
<b>Quebec</b>	Unknown	N/A	N/A			(blank)

*Column options*

**Present?:** Yes; No; Unknown; No data; (blank) or Choose an Item

**Abundance and Distribution:** Declining; Increasing; Stable; Unknown; Extirpated; N/A; (blank) or Choose an item

**SGCN?:** Yes; No; Unknown; (blank) or Choose an item

### **Monitoring in New York** *(specify any monitoring activities or regular surveys that are conducted in New York):*


As part of a State Wildlife Grant, NYSDEC Region 8 Fish and Wildlife staff conducted a native freshwater mussel baseline inventory of tributaries in central and western New York, 2009 to 2020.

### **Trends Discussion** *(insert map of North American/regional distribution and status):*

Trends for New York populations are difficult to determine as most historic data comes from opportunistic naturalist collections, as opposed to more comprehensive baseline surveys. For example, mussels were documented for the first time in 50 of the 106 streams surveyed to date by the Southern Lake Ontario mussel inventory project (Mahar & Landry 2013). This is because many of these streams had never before been surveyed for mussels, not because mussel distribution has dramatically increased. In North America, approximately 2/3 to ¾ of native mussel species are extinct, listed as endangered or threatened, or are in need of conservation status (Williams et al., 1993; Stein et al., 2000). Based on New York's Natural Heritage S-rank, sparse historical data, and the plight of North America's freshwater mussels, it is assumed that trends are declining due to a myriad of environmental stressors.




Figure 1. Mucket distribution (IUCN Redlist 2024)



**Figure 2.** Mucket status (NatureServe 2024)

**III. New York Rarity** (provide map, numbers, and percent of state occupied)



**Figure 3.** Records of mucket in New York (NYSDEC 2024)

	<b># of Distinct Waterbodies</b>	<b>% of State</b>
<b>Total</b>	<u>10</u>	<u>1.2%</u>

**Table 1.** Records of mucket in New York.

**Details of historic and current occurrence:**

2024: *O. ligamentina* has been found in 21 of 1802 HUC 12 watersheds (1.2%) and 10 waterbodies in New York.

Historically, this species was widespread and abundant in the Allegheny basin. There are also old records from the Niagara River, Oak Orchard Creek, and possibly Tonawanda Creek (Strayer & Jirka 1997).

Since 1970, *O. ligamentina* has been found in ten New York State waterbodies.

*O. ligamentina* is currently found in medium and large creeks and rivers of the greater Allegheny River system, in which it is arguably the most abundant species (Strayer & Jirka 1997; Smith & Crabtree 2005; Smith & Meyer 2008; Smith and Crabtree 2009; The Nature Conservancy 2009), and often is the dominant species in medium-gradient habitats (such as the main stem of the Allegheny River) (The Nature Conservancy 2009). The Nature Conservancy found 4,163 live *O. ligamentina* at 61 of 105 survey sites with greatest total catches (up to 117 per hour) in the Allegheny River. *O. ligamentina* populations were found throughout both the Upper Allegheny and Conewango sub-basins, including Oswayo Creek, Olean Creek, Conewango Creek, Cassadaga Creek, and the mainstem of the Allegheny River. Populations were considered viable at 82% of the sites where they were found (The Nature Conservancy 2009).

Between 1988 and 1990, live *O. ligamentina* were found at several locations in the New York portion of French Creek. During this same time period, old shells were found in Tonawanda Creek in the Erie Basin (New York Natural Heritage Program 2013).

In 2013 over 50 live *O. ligamentina* were found in French Creek, southwest of French Creek town, Chautauqua County (Burlakova, Karatayev, unpublished data).

**New York’s Contribution to Species North American Range:**

Percent of North American Range in NY	Classification of NY Range	Distance to core population, if not in NY
1-25%	Peripheral	230 miles

Column options

Percent of North American Range in NY: 100% (endemic); 76-99%; 51-75%; 26-50%; 1-25%; 0%; Choose an item

Classification of NY Range: Core; Peripheral; Disjunct; (blank) or Choose an item

**IV. Primary Habitat or Community Type** (from NY crosswalk of NE Aquatic, Marine, or Terrestrial Habitat Classification Systems):

- a. **Size/Waterbody Type:** Small River to Medium River
- b. **Geology:** Moderately Buffered
- c. **Temperature:** Transitional Cool to Warm
- d. **Gradient:** Low Gradient to Moderate-High Gradient

**Habitat or Community Type Trend in New York**

Breeder in NY?	Non-breeder in NY?	Migratory Only?	Summer Resident?	Winter Resident?	Anadromous/Catadromous?
Yes	No	No	Yes	Yes	(blank)

Column options

Habitat Specialist and Indicator Species: Yes; No; Unknown; (blank) or Choose an item

Habitat/Community Trend: Declining; Stable; Increasing; Unknown; (blank) or Choose an item

**Habitat Discussion:**

Mucket typically inhabits large creeks and rivers (Strayer & Jirka 1997; Cummings & Mayer 2002; NatureServe 2013). It is "best fitted for the rough parts, riffles with strong current and heavy gravel

and rocks," but may also be found in sandy mud or gravel along stream margins (Ortmann, 1919). This early habitat assessment is consistent with Watters et al. (2009) and Strayer & Jirka (1997), who note that the species is most common in cobble and sand in moving water, although habitats in New York range from stony riffles to soft-bottomed pools. It may rarely occur in shallow water areas of large lakes (NatureServe 2013).

## V. Species Demographic, and Life History:

Breeder in NY?	Non-breeder in NY?	Migratory Only?	Summer Resident?	Winter Resident?	Anadromous/Catadromous?
Yes	No	No	Yes	Yes	(blank)

Column options

First 5 fields: Yes; No; Unknown; (blank) or Choose an item

Anadromous/Catadromous: Anadromous; Catadromous; (blank) or Choose an item

**Species Demographics and Life History Discussion** (include information about species life span, reproductive longevity, reproductive capacity, age to maturity, and ability to disperse and colonize):

Upstream males release sperm into the water. Females downstream take up the sperm with incoming water. Fertilization success may be related to population density, with a threshold density required for any reproductive success to occur. Eggs are fertilized within the female. Like nearly all North American mussels, *O. ligamentina* must parasitize an often-specific vertebrate host to complete its life cycle. It is suspected that some mussel populations are not recruiting because their hosts no longer occur with them. Once released by the female, glochidia must acquire a suitable host or die, usually within 24-48 hours. After attaching to a suitable host, glochidia encyst, usually at the fish's gills or fins and receive food and dispersal. Once the glochidia metamorphose into juveniles, they drop from the host. If they land in suitable habitat, they will burrow into the substrate, where they may remain for several years (Watters et al. 2009). Less than one in a million glochidia survive to adulthood (Young & Williams, 1984).

In the adult form, freshwater mussels are basically sessile; movement is limited to a few meters of the lake or river bottom. The only time that significant dispersal can take place is during the parasitic phase. Infected host fishes can transport the larval unionids into new habitats, and can replenish depleted populations with new individuals. Dispersal is particularly important for genetic exchange between populations. Dispersal is likely a slow process for mussels which use resident fishes with limited home ranges as their hosts (COSEWIC as cited in NatureServe 2013).

It has an equilibrium life history strategy, characterized primarily by long life span, mostly short term brooding, low to moderate growth rate, and late maturity, with low reproductive effort and fecundity that increases slowly after maturation. This life history strategy is considered to be favored in stable, productive habitats (Haag 2012).

*O. ligamentina* reaches sexual maturity at 4 to 6 years and can live up to 25 years. This species is bradyctictic, with eggs developing in mid-summer and glochidia are present in the female from September to the following May to August (Watters et al. 2009). *O. ligamentina* has been known to use many species of warm water fish as hosts and glochidia have been found to transform on rock bass (*Ambloplites rupestris*), central stoneroller (*Campostoma anomalum*), silverjaw minnow (*Notropis buccatus*), banded killifish (*Fundulus diaphanus*), green sunfish (*Lepomis cyanellus*), smallmouth bass (*Micropterus dolomieu*), largemouth bass (*Micropterus salmoides*), white bass (*Morone chrysops*), yellow perch (*Perca flavescens*), white crappie (*Pomoxis annularis*), and black

crappie (*Pomoxis nigromaculatus*). It is able to infest American eel (*Anguilla rostrata*), common carp (*Cyprinus carpio*), bluegill (*Lepomis macrochirus*), tadpole madtom (*Noturus gyrinus*), and sauger (*Sander canadensis*) (Watters et al. 2009; Strayer & Jirka 1997).

## **VI. Threats** (from NY 2015 SWAP or newly described):

### **Agricultural Runoff**

New York's largest populations of *O. ligamentina* are found in the Allegheny River upstream of Olean, with additional viable populations between Olean and Salamanca (The Nature Conservancy 2009). Roughly half of these sections of stream are bordered by agriculture, primarily in the Olean/Allegheny area. Although the mid reaches of Cassadaga Creek are quite forested, both the upstream portions of Cassadaga Creek and the lower portions of Conewango Creek, in which *O. ligamentina* have been found, are influenced by limited agriculture. The French Creek watershed is also highly agricultural (New York State Landcover 2010). Aquatic habitat lacking vegetated buffers of adequate width are threatened by runoff from urban areas, roads, lawns, and agricultural land (Gillis 2012). If best management practices are not closely adhered to, mussel habitat adjacent to wood harvest or agricultural land is subjected to pesticide, fertilizer, and silt/sediment runoff. During recent mussel surveys in Western and Central New York, it has been documented that sufficient vegetated riparian buffers are often lacking along known mussel streams (Mahar & Landry 2013), indicating that runoff is a major threat to resident mussel populations.

The presence of pesticides and fertilizers in our rural watersheds is nearly ubiquitous (Haag 2012). And because pesticides and their associated surfactants adsorb onto sediment particles, sedimentation may act as a vector for their transport into the aquatic system (Haag 2012). Mussels are more sensitive to pesticides than many other animals (Watters et al. 2009). Although effects of pesticides are species-specific, sub-lethal levels of PCBs, DDT, malathion, and other compounds inhibit respiratory efficiency and accumulate in the tissues. Atrazine and permethrin at sublethal concentrations reduced juvenile growth (Bringolf et al. 2007a, 2007b) and environmental levels of atrazine altered mussel movement and aggregation behavior (Flynn and Spellman 2009). Pesticides can affect mussels in many ways, but the full range of long-term effects remains unknown (Haag 2012).

Fertilizer runoff is also a concern. High inputs of nitrogen from fertilizers can cause increases in ammonia in the water and the substrate, leading to direct toxicity for a wide range of mussel species. Mussels, especially in their early life stages, are more sensitive to un-ionized ammonia than other organisms, and high sensitivity is seen across a range of species and life histories (Haag 2012). In addition, ammonia adsorbs to sediment particles, resulting in higher nitrogen concentrations in the substrate than in the overlying water. The nitrogen present in the interstitial spaces in the substrate is thought to result in juvenile mortality and to prevent recruitment by some mussel species (Strayer and Malcom 2012). Studies have suggested decreasing sediment loads entering aquatic systems as the best way to decrease the impact of numerous stressors for mussels in general (Roley et al. 2012).

### **Treated Wastewater**

The habitat of *O. ligamentina* receives treated wastewater from the cities of Olean, Salamanca, and the village of Portville, either directly to the Allegheny River or through tributaries. Cassadaga Creek receives treated effluent from the city of Jamestown sewage treatment plant (SPDES 2007). Recent studies show that mussel richness and abundance decrease with increased proximity to sewage effluent (Wildenberg 2012). The input of biomaterial from wastewater treatment plants

depletes dissolved oxygen levels, negatively impacting mussels. Ammonia from wastewater treatment plants has been found to be toxic to glochidia (Goudraeu et al. 1993) and at sub-lethal exposure, adult mussels exhibit decreased respiratory efficiency (Anderson et al. 1978). Endocrine disrupters from pharmaceuticals are also present in municipal sewage effluents and are increasing common in rivers and lakes (Haag 2012). In mussels, chronic exposure to estrogenic compounds in effluents caused feminization of male mussels, but these individuals did not produce eggs, suggesting major disruption of reproductive function (Gagne et al. 2011). The long-term effects of these compounds on mussels are unknown (Haag 2012). It should be noted that in the Susquehanna Basin, Harman and Lord (2010) found no evidence that wastewater treatment plants were responsible for reductions in mussel species of greatest conservation need.

### **Runoff from Developed Land**

The habitat of *O. ligamentina* receives storm water runoff from the cities of Olean, Salamanca, and the village of Portville, either directly to the Allegheny River or through tributaries. Cassadaga Creek receives Jamestown's urban runoff via the Chadokoin River. All ten New York waterbodies that host *O. ligamentina* populations are intermittently bordered by interstate highways, state routes, and/or local roads (New York State Landcover, 2010). These developed lands are likely sources runoff containing metals and road salts. Mussels are particularly sensitive to heavy metals, more so than many other animals used in toxicological tests (Keller & Zam 1991). Low levels of metals may interfere with the ability of glochidia to attach to the host (Huebner & Pynnonen 1992), suggesting that U.S. EPA ambient water quality criteria may not adequately protect mussels from toxic metals (Wang et al. 2011). In addition, increases in salinity from the runoff of salt used for clearing roads in winter or output from cheese production may be lethal to glochidia and juvenile mussels (Keller & Zam 1991; Liquori & Insler 1985; Pandolfo et al. 2012). Based on these studies, the U.S. EPA's ambient water quality criterion for acute chloride exposures may not be protective of all freshwater mussels (Pandolfo et al. 2012).

### **Flood Control Projects**

Large stretches of *O. ligamentina* habitat are within the leveed portions of the Allegheny River, Olean Creek, and Oswayo Creek ("New York State Flood Protection" 2013). These structures confine larger rivers, preventing the river from inundating its natural floodplains and wetlands to minimize flood damage. Additionally, many smaller streams have been channelized and bermed by landowners and highway departments to protect farm fields and other structures. Channelization and dredging associated with flood control projects are catastrophic to mussels and have been implicated in the decline of some populations (Watters et al. 2009). The result of these projects is altered seasonality of flow and temperature regimes, increased stream velocities, unstable substrates, changed patterns of sediment scour and deposition, including streambank erosion, altered transport of particulate organic matter (the food base for mussels), and a general degradation of stream habitat (Benke 1999; Yeager 1993; Nedeau 2008).

### **Other Habitat Modifications**

Ecosystem modifications, such as isolated occurrences of flood control channel dredging, instream work associated with bridge replacement, or gravel mining, and vegetation removal kill mussels and destroy their habitat. For example, dredging for vegetation removal has been shown to remove up to 23% of mussels in spoils (Aldridge 2000). Further evidence for disruption was provided by mussel surveys adjacent to approximately 20 river miles of Conewango Creek that had been channelized and straightened in the first half of the 20th century. The resulting "dredge" had no riffle or run habitat and sites just below and above this channelized section contained few or no mussels (The Nature Conservancy 2009). Although limited in geographic scope these habitat modification activities have long term impacts on mussels and their distribution (Aldridge 2000).

### **Invasive Species**

Invasive zebra mussels (*Dreissena polymorpha*) pose a potential threat to *O. ligamentina* populations in Cassadaga and Conewango Creeks, where they are present in the lower reaches. Chautauqua Lake's connection to Cassadaga Creek, Chadakoin River, is the main source of this exotic invasive (The Nature Conservancy 2009), that has been repeatedly cited as a threat to native mussel populations (Strayer & Jirka 1997; Watters et al. 2009). Studies have shown that *O. ligamentina* are significantly stressed by zebra mussels (Baker & Hornbach 1997). En masse, Dreissenids outcompete native mussels by efficiently filtering food and oxygen from the water. They reduce reproductive success by filtering native mussel male gametes from the water column and they can foul the shells of the native mussels to the point that their valves can no longer open. In heavily invested areas, they may transform a habitat by hardening the substrate, such that dislodged mussels are not able to rebury (USFWS 1994). Although zebra mussels will continue to cause problems for Chautauqua Lake, they currently appear to have minimal impact downstream. However, precautions should be taken to avoid invasions by zebra mussels to upstream locations, especially the headwater lakes in the Cassadaga system. Monitoring for zebra mussels in these lakes may provide early detection of this invader (The Nature Conservancy 2009).

### **Impoundments**

Across its range, impoundments likely contributed to the reduced distribution of mussels that we see today. Vaughn and Taylor (1999) observed a mussel extinction gradient with a gradual, linear increase in mussel species richness and abundance with increasing distance downstream from impoundments. Species and their hosts that require shallow, oxygenated, fast-flowing water quickly are eliminated. Continuously cold water from both increased water depth upstream of the dam and dam discharges downstream of the dam may prevent reproduction. Impoundment increases silt load and eutrophication, resulting in changes in the fish fauna, and therefore the availability of hosts. Dams represent distributional barriers to fish hosts, and therefore to the mussels themselves. The zoogeographic patterns of several species suggest a dam-limited range. Dams also act as sediment traps, often having many feet of silt and debris caught on their upstream side. These areas generally are without mussels. Below the dam, the tailwaters often have dense mussel beds, as these reaches are the only areas left that still have oxygenated, fast moving water. This is exemplified by the distribution of beds in the lower Muskingum River, Ohio (Stansbery & King 1983; ESI 1993c).

In addition, improperly sized and poorly installed or poorly maintained culverts have impacts similar to dams in that they fragment habitat, preventing the movement by host fish, and effectively isolating mussel populations. And because culverts are located at nearly every road-stream intersection, there is the potential for landscape level fragmentation of mussel habitat.

<b>Threat Level 1</b>	<b>Threat Level 2</b>	<b>Threat Level 3</b>	<b>Spatial Extent</b>	<b>Severity</b>	<b>Immediacy</b>	<b>Trend</b>	<b>Certainty</b>
1. Residential and Commercial	1.3 Tourism & Recreation Areas	1.3.5 Docks & marinas	Restricted	Serious	Immediate	Stable and ongoing	High
3. Energy Production & Mining	3.3 Renewable Energy	3.3.1 Hydroelectric dams	Large	Moderate	Near-term	Intensifying	High
4. Transportation & Service Corridors	4.1 Roads & Railroads	4.1.3 Bridges	Restricted	Serious	Long-term	Stable and ongoing	High
6. Human Intrusions & Disturbance	6.3 Work & Other Activities	6.3.1 Research activities	Restricted	Slight	Immediate	Stable and ongoing	Low
7. Natural System Modifications	7.2 Dams & Water Management/Use	7.2.1 Water level management using dams	Large	Moderate	Near-term	Intensifying	High
7. Natural System Modifications	7.2 Dams & Water Management/Use	7.2.6 Withdrawl of surface water	Restricted	Moderate	Long-term	Unknown	High
7. Natural System Modifications	7.2 Dams & Water Management/Use	7.2.3 Water management using culverts	Large	Slight	Long-term	Unknown	High
7. Natural System Modifications	7.3 Other Ecosystem Modifications	7.3.3 Natural erosion & sedimentation	Large	Moderate	Near-term	Intensifying	High
8. Invasive & Other Problematic Species	8.1 Invasive Non-Native Plants & Animals	8.1.3 Aquatic animals	Large	Extreme	Near-term	Stable and ongoing	High
8. Invasive & Other Problematic Species	8.2 Problematic Native Plants & Animals	8.2.5 Increased predation by mesopredators	Restricted	Slight	Long-term	Intensifying	Moderate
8. Invasive & Other Problematic Species	8.5 Intrinsic Biological Limitations	8.5.1 Loss of genetic diversity	Large	Moderate	Long-term	Intensifying	High
8. Invasive & Other Problematic Species	8.5 Intrinsic Biological Limitations	8.5.2 Depends on another species that has declined	Restricted	Moderate	Long-term	Unknown	High
9. Pollution	9.1 Domestic & Urban Wastewater	9.1.1 Domestic wastewater	Small	Moderate	Near-term	Unknown	High

9. Pollution	9.1 Domestic & Urban Wastewater	9.1.2 Runoff	Restricted	Moderate	Near-term	Unknown	Moderate
9. Pollution	9.3 Agricultural & Forestry Effluents	9.3.2 Soil erosion, sedimentation	Large	Extreme	Near-term	Unknown	High
9. Pollution	9.3 Agricultural & Forestry Effluents	9.3.3 Herbicides & pesticides	Restricted	Moderate	Near-term	Unknown	Moderate
11. Climate Change	11.3 Changes in Temperature Regimes	11.3.3 Gradual temperature change	Large	Moderate	Long-term	Intensifying	Low
11. Climate Change	11.3 Changes in Temperature Regimes	11.3.4 Increase in temperature fluctuations	Large	Moderate	Long-term	Intensifying	Low
11. Climate Change	11.4 Changes in Precipitation & Hydrological Regimes	11.4.2 Droughts	Restricted	Moderate	Long-term	Intensifying	High
11. Climate Change	11.5 Storms & Severe Weather	11.5.1 Storms & severe weather	Restricted	Moderate	Long-term	Intensifying	Moderate

**Table 2.** Threats to Mucket.

**Are there regulatory mechanisms that protect the species or its habitat in New York?**

Yes:

No:

Unknown:

**If yes, describe mechanism and whether adequate to protect species/habitat:**

Mussel habitats receive some generic protection under several New York State regulations (NYCRR) promulgated under the authority of the New York Environmental Conservation Law (ECL), specifically Part 608 of the NYCRR: Use and Protection of Waters, and Part 617 of the NYCRR: State Environmental Quality Review (SEQR). Part 608 provides protection of some mussel habitats by regulating and requiring environmental review of the modification or disturbance of any “protected stream”, its bed or bank, and removal of sand, gravel or other material from its bed or banks (608.2 Disturbance of Protected Streams). This does not provide adequate protection of mussels and their habitats as it only protects streams or particular portions of a streams for which there has been adopted by NYSDEC or any of its predecessors any of the following classifications or standards: AA, AA(t), A, A(t), B, B(t) C(t), or Streams designated (t)(trout) also include those more specifically designated (ts)(trout spawning). Mussel habitats may also receive some additional protections as the construction, repair, breach or removals of dams, and the excavation and placement of fill in navigable waters are subject to regulation and environmental review under Part 608, 608.3 and 608.5 respectively. Under Part 608, projects requiring a permit can be conditioned by NYSDEC to include best management practices, such as sediment and erosion protections. Through the review process, these projects can also be modified to reduce impacts in order to meet permit issuance standards.

Under Part 608, protection of unlisted species of mussels is general and relatively limited. More importantly, Class C and D waters with mussels do not receive protection under these regulations. A significant portion of the New York’s mussel resources occur within Class C and D waters. An additional but not insignificant gap in protection occurs because agricultural activities consisting of the crossing and re-crossing of a protected stream by livestock or wheeled farming equipment normally used for traditional agricultural purposes or of withdrawing irrigation water in a manner which does not otherwise alter the stream, are exempt from these regulations and environmental review.

Water quality certifications required by Section 401 of the Federal Water Pollution Control Act, Title 33 United States Code 1341(see subdivision (c)of this Section) may provide protection for freshwater mussels and their habitats from some activities that would potentially have adverse impacts by regulating construction or operation of facilities that may result in any discharge into navigable waters. Water quality certifications set water quality-related effluent limitations, water quality standards, thermal discharge criteria, effluent prohibitions and pretreatment standards for projects on navigable waters.

The State Environmental Quality Review (SEQR, Part 617 NYCRR) may also protect mussels and their habitats by requiring the consideration of environmental factors into the existing planning, review and decision-making processes of state, regional and local government agencies for activities that require discretionary approval. SEQR requires the preparation of an Environmental Impact Statement, including an alternatives analysis, for those activities that may result in a substantial adverse change in ground or surface water quality; a substantial increase in potential for erosion, flooding, leaching or drainage problems; the removal or destruction of large quantities of vegetation or fauna; substantial interference with the movement of any resident or migratory fish or wildlife species; impacts on a significant habitat area; substantial adverse impacts on a

threatened or endangered species of animal or plant, or the habitat of such a species; other significant adverse impacts to natural resources; or, a substantial change in the use, or intensity of use, of land including agricultural, open space or recreational resources, or in its capacity to support existing uses.

New York State has numerous laws and regulations that both directly or indirectly protect waters of the state (mussel habitats) including regulations governing direct discharges to surface and groundwater, storm water, agricultural activities, pesticides, flood control, and dams. Without these regulations, mussels would certainly be in worse shape; however, most of these generic protections are not adequate in scope or specific enough to mussel threats to protect the mussel resources of New York State.

**Describe knowledge of management/conservation actions that are needed for recovery/conservation, or to eliminate, minimize, or compensate for the identified threats:**

- Priority conservation efforts for this species should focus on, but not be limited to, the Allegheny River upstream of Olean.
- Through landowner incentive programs or regulation, riparian buffers, particularly those that also provide shade, should be added/maintained/widened, along agricultural fields, subdivisions, and along major roads to decrease the levels of nitrogen, pesticides, sediment, heavy metals, and salts from entering these aquatic systems, as well as to moderate water temperature. Studies have suggested decreasing sediment loads entering aquatic systems as the best way to decrease the impact of numerous stressors for mussels in general (Roley & Tank 2012).
- Require all state agencies to maintain appropriate vegetative buffers along streams, rivers and lakes on state-owned or state managed properties.
- Develop and implement a comprehensive monitoring strategy that identifies protocols, including locations and specific intervals, for regular monitoring of known mussel populations to detect assess trends and detect dangerous declines.
- Mussel sensitivity to particular pollutants should be considered or addressed in the regulation of wastewater and stormwater discharges to groundwater and surface waters, State Pollutant Discharge Elimination Systems (SPDES). This should be reflected in effluent limitations for discharges, including discharges from P/C/I facilities (Private/Commercial/Industrial), CAFO facilities (Concentrated Animal Feeding Operations), High Volume Hydraulic Fracturing Discharges, and Wastewater treatment plants, etc. Discharges whose receiving waters have mussels, particularly those with known populations of mussels listed as Endangered, Threatened, Special concern or SGCN, should be carefully reviewed for potential impacts to mussels. For example, deleterious levels of ammonia (a component of many types of discharges) and molluscicides (a commonly used water treatment chemical in discharged water) should not be permitted.
- Establish a protocol whereas DEC staff work closely with Flood Control Management to reduce impacts to native mussels during maintenance flood control projects.
- Coordinate with local wastewater treatment facilities to improve ammonia removal of treated discharge. This has been documented as a threat to Unionids at multiple life stages, and therefore needs to be addressed (Gillis 2012).

- Establish a protocol whereas DEC staff work closely with state and local highway departments to reduce impacts to native mussels during maintenance and construction projects.
- Replace culverts that disrupt aquatic habitat connectivity to allow for passage of small fish species.
- NYSDEC should consider sensitivity of freshwater mussels to specific pollutants in the establishment and setting of water quality standards and TMDLs for waters containing freshwater mussels. A Total Maximum Daily Load (TMDL) specifies the maximum amount of a pollutant that a waterbody can receive and still meet water quality standards. TMDLs account for all contributing sources (e.g. point & nonpoint sources, and natural background levels), seasonal variations in the pollutant load, and incorporate a margin of safety that accounts for unknown or unexpected sources of the pollutant. In essence, a TMDL defines the capacity of the waterbody to absorb a pollutant and still meet water quality standards. The Clean Water Act requires states to identify waterbodies that do not meet water quality standards after application of technology-based effluent limitations. For these "impaired waters," states must consider the development of alternative strategies, including TMDLs, for reducing the pollutants responsible for the failure to meet water quality standards.

Action Category	Action	Description
A.1 Direct Habitat Management	A.1.2.4.1 Plant for erosion management	Implement riparian buffers to reduce sedimentation
A.1 Direct Habitat Management	A.1.3.0.0 Mitigate human environmental impact	Mitigations should be focused on the Allegheny River upstream of Olean, NY.
A.1 Direct Habitat Management	A.1.3.3.0 Remove and improve anthropogenic infrastructure	Replace culverts that disrupt habitat connectivity and fish passage
B.3 Outreach	B.3.1.3.0 Targeted Communication	-Coordinate with local wastewater treatment to improve ammonia removal in treated discharge  -Work with Highway Departments to reduce impacts on mussels
C.6 Design and Plan Conservation	C.6.5.1.2 Develop a strategy, guideline, monitoring plan or follow-up in a protected area	Develop a plan to monitor mussel population trends

Action Category	Action	Description
C.7 Legislative and Regulatory Framework or Tools	C.7.1.3.0 Create, amend, or influence regulation	Require state agencies to have appropriate vegetative buffers alongside water on state land
C.7 Legislative and Regulatory Framework or Tools	C.7.2.3.0 Create or amend standards	Implement TMDLs to meet water quality standards.
C.8 Research and Monitoring	C.8.2.1.2 Monitoring and evaluating the results of project activities	

**Table 3.** Recommended conservation actions for mucket.

The Comprehensive Wildlife Conservation Strategy (NYSDEC 2006) includes recommendations for the following actions for freshwater mussels:

**Habitat management:**

- Manage areas of important mussel populations by controlling degradation factors (e.g.. Controlling livestock access, point source or non-point source pollution, flow alteration, etc.)
- Develop methods to improve and restore freshwater bivalve habitat.

**Habitat research:**

- Conduct research to determine habitat parameters necessary for good populations of each species of species-at-risk listed mussels.
- Research flow requirements of freshwater bivalves and model the effects of flow changes both in volume and timing.
- Research all parameters of mussel habitat requirements including temperature, substrate, fish, flow, food, etc.

**Habitat restoration:**

- Restore degraded habitat areas to allow for recolonization or reintroduction of listed mussels.

**Invasive species control:**

- Develop a monitoring/control plan that includes measures to detect invasive species problematic to freshwater bivalves in all New York watersheds and actions that will be taken to control them before they become threats.
- Conduct research on control of exotic bivalve species that compete with native mussels and exotic crustaceans or fish which may prey on them.

**Life history research:**

- Research effects of pesticides and other chemicals, including ammonia, on all life stages of freshwater bivalves: sperm/egg, glochidia, larva, adults.
- Determine fish hosts for species where this is not known for populations living in New York.
- Research population dynamics of listed mussel species including connectivity of populations or subpopulations and genetic distinctness of populations or subpopulations.

- Determine or confirm breeding phenology and habitat conditions necessary for successful breeding for listed mussels (e.g., mussel density, pop. level of fish host, temp, flow).

**New regulation:**

- Ban the importation of fish that feed on freshwater mollusks (e.g. black carp).
- Require inclusion of all stages of freshwater mussels in testing for approval of new pesticides in New York.

**Other action:**

- Develop an outreach program to private landowners through the Landowner Incentive Program to educate the public about freshwater mussel protection and initiate projects to prevent or repair impacts from land use on mussels.
- Increase regional permit control of development and highway projects that may impact native mussels.
- Develop standard monitoring/survey protocols for development projects in all watersheds in New York.
- Evaluate threats to mussels in each New York watershed and prioritize areas for actions to address the threats.
- Research the best survey methods both for detection of rare species and evaluation of population status and trends.

**Population monitoring:**

- Conduct population estimates of species-at-risk listed mussel species in NY
- Conduct surveys to determine distribution of species-at-risk listed mussel species in NY.

**Regional management plan:**

- Incorporate freshwater mussel goals and objectives into regional water quality and fish management plans and policies.

**Relocation/reintroduction:**

- Where appropriate, reintroduce listed mussels into appropriate habitat within their historic range.

**Statewide management plan:**

- Incorporate freshwater mussel goals and objectives into statewide water quality and fish management plans and policies.

## VII. References

Aldridge, D. C. (2000). The impacts of dredging and weed cutting on a population of freshwater mussels (Bivalvia: Unionidae). *Biological Conservation*, 95(3), 247-257.

Anderson, K. B., Sparks, R. E., & Paparo, A. A. (1978). Rapid assessment of water quality, using the fingernail clam, *Musculium transversum*: Final Report. University of Illinois, Urbana. 130p.

- Baker, S. M., & Hornbach, D. J. (1997). Acute physiological effects of zebra mussel (*Dreissena polymorpha*) infestation on two unionid mussels, *Actiononaias ligamentina* and *Amblema plicata*. *Canadian Journal of Fisheries and Aquatic Sciences*, 54(3), 512-519.
- Benke, A.C. (1990). A perspective on America's vanishing streams. *Journal of the N. American Benthological Society*: 9: 77-88.
- Bringolf, R. B., Cope, W. G., Eads, C. B., Lazaro, P. R., Barnhart, M. C., & Shea, D. (2007). Acute and chronic toxicity of technical-grade pesticides to glochidia and juveniles of freshwater mussels (unionidae). *Environmental Toxicology and Chemistry*, 26(10), 2086-2093.
- Bringolf, R. B., Cope, W. G., Barnhart, M. C., Mosher, S., Lazaro, P. R., & Shea, D. (2007). Acute and chronic toxicity of pesticide formulations (atrazine, chlorpyrifos, and permethrin) to glochidia and juveniles of *Lampsilis siliquoidea*. *Environmental Toxicology and Chemistry*, 26(10), 2101-2107.
- COSEWIC. (2003). COSEWIC assessment and status report on the kidneyshell *Ptychobranhus fasciolaris* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa, Canada.
- Cummings, K. S., & Mayer, C. A. (1992). Field guide to freshwater mussels of the Midwest (p. 194). Champaign, Illinois: Illinois Natural History Survey.
- Flynn, K., & Spellman, T. (2009). Environmental levels of atrazine decrease spatial aggregation in the freshwater mussel, *Elliptio complanata*. *Ecotoxicology and Environmental Safety*, 72(4), 1228-1233.
- Graf, D. and K. Cummings. (2011). MUSSELp Evolution: North American Freshwater Mussels. The MUSSEL Project. The University of Wisconsin. [http://mussel-project.uwsp.edu/evol/intro/north\\_america.html](http://mussel-project.uwsp.edu/evol/intro/north_america.html)
- Graf, Daniel & Cummings, Kevin. (2021). A 'big data' approach to global freshwater mussel diversity (Bivalvia: Unionoida), with an updated checklist of genera and species. *Journal of Molluscan Studies*. 87. 10.1093/mollus/eyaa034.
- Gagné, F., Bouchard, B., André, C., Farcy, E., & Fournier, M. (2011). Evidence of feminization in wild *Elliptio complanata* mussels in the receiving waters downstream of a municipal effluent outfall. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 153(1), 99-106.
- Gillis, P. L. (2012). Cumulative impacts of urban runoff and municipal wastewater effluents on wild freshwater mussels (*Lasmigona costata*). *Science of the Total Environment*, 431, 348-356.
- Goudraeu, S. E., Neves, R. J., & Sheehan, R. J. (1993). Effects of wastewater treatment plant effluents on freshwater mollusks in the upper Clinch River, Virginia, USA. *Hydrobiologia*, 252(3), 211-230.
- Haag, W. R. (2012). *North American freshwater mussels: natural history, ecology, and conservation*. Cambridge University Press.
- Harman, W.N. and P.H. Lord (2010). Susquehanna Freshwater Mussel Surveys, 2008-2010. Final report submitted to New York State Department of Environmental Conservation. SUNY Oneonta. Cooperstown, NY. 24 pp, plus appendix.

- Huebner, J. D., & Pynnönen, K. S. (1992). Viability of glochidia of two species of *Anodonta* exposed to low pH and selected metals. *Canadian Journal of Zoology*, 70(12), 2348-2355.
- Keller, A. E., & Zam, S. G. (1991). The acute toxicity of selected metals to the freshwater mussel, *Anodonta imbecilis*. *Environmental Toxicology and Chemistry*, 10(4), 539-546.
- Liquori, V. M., & Insler, G. D. (1985). Gill parasites of the white perch: Phenologies in the lower Hudson River. *New York Fish and Game Journal*, 32(1), 71-76.
- Mahar, A.M. and J.A. Landry. (2014). State Wildlife Grants Final Report: Inventory of Freshwater Mussels in New York's Southeast and Southwest Lake Ontario Basins, 2008-2013. New York State Department of Environmental Conservation. Avon, NY.
- Mahar, A.M. and J.A. Landry. 2013. NYSDEC SWAP 2015 Species Status Assessment for Mucket. Prepared on June 2013. Revised by Samantha Hoff on February 25, 2014.
- MolluscaBase eds. (2024). MolluscaBase. *Ortmanniana ligamentina* (Lamarck, 1819). Accessed at: <https://www.molluscabase.org/aphia.php?p=taxdetails&id=857466> on 2024-10-31
- Natural Heritage Program Element Occurrences [ARC/INFO coverages] (2013). New York Natural Heritage Program, Albany, NY. Available: NYS Department of Environmental Conservation Master Habitat Data Bank's Data Selector.
- NatureServe. 2013. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: February 12, 2013).
- Nedeau, E.J. 2008. *Freshwater Mussels and the Connecticut River Watershed*. Connecticut River Watershed Council, Greenfield, Massachusetts. xviii+ 132 pp.
- New York State Department of Environmental Conservation. (2006). *New York State Comprehensive Wildlife Conservation Strategy*. Albany, NY: New York State Department of Environmental Conservation.
- New York State Flood Protection Project Details and Maps (2013). Retrieved from Department of Environmental Conservation website: <http://www.dec.ny.gov/lands/59934.html>
- New York State Landcover, Version 1. [SDE raster digital data] (2010). National Gap Analysis Program. Moscow, Idaho. Available: NYS Department of Environmental Conservation Master Habitat Data Bank's Data Selector.
- Northeastern Aquatic Habitat Classification System (NAHCS) GIS map for streams and rivers, [vector digital data] (2010). US Environmental Protection Agency, the US Geological Survey, and The Nature Conservancy Eastern Conservation Science. Boston, MA. Available: NYS Department of Environmental Conservation Master Habitat Data Bank's Data Selector.
- Ortmann, A. E. (1919). *Monograph of the Naiades of Pennsylvania*. (Vol. 8, No. 1). Board of Trustees of the Carnegie Institute.
- Pandolfo, T. J., Cope, W. G., Young, G. B., Jones, J. W., Hua, D., & Lingenfelter, S. F. (2012). Acute effects of road salts and associated cyanide compounds on the early life stages of the unionid mussel *Villosa iris*. *Environmental Toxicology and Chemistry*, 31(8), 1801-1806.

- Roley, S.S. 2012. The influence of floodplain restoration on stream ecosystem function in an agricultural landscape. (unpublished doctoral dissertation). University of Notre Dame, Notre Dame, Indiana. Submitted for publishing with Tank, J.L.
- Roley, S. S., J. Tank, and M. A. Williams (2012), Hydrologic connectivity increases denitrification in the hyporheic zone and restored floodplains of an agricultural stream, *J. Geophys. Res.*
- Smith, T. A. and D. Crabtree. 2005. Freshwater Mussel (Unionidae) and Fish Assemblage Habitat Use and Spatial Distributions in the French Creek Watershed: Reference for Western Pennsylvania Unionid Protection and Restoration. Unpublished July 30, 2005 Final Report submitted to the Pennsylvania Fish and Boat Commission iv + 180pp.
- Smith, T.A., and D.L. Crabtree. (2009). Freshwater mussel (Unionidae: Bivalvia) distributions and densities in French Creek, Pennsylvania. *Northeastern Naturalist*.
- Smith, T. A. and E. Meyer. (2008). Freshwater Mussel (Unionidae) Surveys in New York Portion of the Allegheny River Watershed, Year Three Report: 2007. February 26. Submitted to The Nature Conservancy. 27pp+app.
- Stansbery, D. H., & King, C. C. (1983). Management of Muskingum River mussel (unionid mollusk) populations. Final Report to the U.S. Department of Commerce, and the Ohio Department of Natural Resources. *Ohio State University Museum of Zoology Reports*. 79 p.
- State Pollutant Discharge Elimination System (SPDES)- New York State [vector digital data]. (2007). Albany, New York: New York State Department of Environmental Conservation. Available: <http://gis.ny.gov/gisdata/inventories/details.cfm?dsid=1010&nysgis=>
- Stein, B. A., Kutner, L. S., Hammerson, G. A., Master, L. L., & Morse, L. E. (2000). State of the states: geographic patterns of diversity, rarity, and endemism. *Precious heritage: the status of biodiversity in the United States*. Oxford University Press, New York, 119-158.
- Strayer, D.L. & K.J. Jirka. (1997). The Pearly Mussels of New York State. *New York State Museum Memoir* (26): 113 pp., 27 pls.
- Strayer, D.L. and Malcom, H.M. 2012. Causes of recruitment failure in freshwater mussel populations in southeastern New York. *Ecological Applications* 22:1780–1790.
- The Nature Conservancy (2009). *Freshwater Mussel (Unionidae) Distributions, Catches, and Measures of their Viability across the Catches, and Measures of their Viability across the Allegheny River Basin in New York*. Report submitted New York State Department of Environmental Conservation. The Nature Conservancy, Central & Western NY Chapter. Rochester, NY. 63 pp.
- U.S. Fish and Wildlife Service. (1994). Clubshell (*Pleurobema clava*) and Northern Riffleshell (*Epioblasma tondosa rangiana*) Recovery Plan. Hadley, Massachusetts. 68 pp.
- Vaughn, C. C. and Taylor, C. M. (1999), Impoundments and the Decline of Freshwater Mussels: a Case Study of an Extinction Gradient. *Conservation Biology*, 13: 912–920.
- Wang, N., Mebane, C. A., Kunz, J. L., Ingersoll, C. G., Brumbaugh, W. G., Santore, R. C., ... & Arnold, W. (2011). Influence of dissolved organic carbon on toxicity of copper to a unionid mussel (*Villosa iris*) and a cladoceran (*Ceriodaphnia dubia*) in acute and chronic water exposures. *Environmental Toxicology and Chemistry*, 30(9), 2115-2125.

- Watters, G. T., Hoggarth, M. A., & Stansbery, D. H. (2009). *The freshwater mussels of Ohio*. Columbus: Ohio State University Press.
- White, E.L., J.J. Schmid, T.G. Howard, M.D. Schlesinger, and A.L. Feldmann. 2011. New York State freshwater conservation blueprint project, phases I and II: Freshwater systems, species, and viability metrics. New York Natural Heritage Program, The Nature Conservancy. Albany, NY. 85 pp. plus appendix.
- Wildenberg, A. (2012, August). Mussel Community Response to Wastewater Effluent in a Midwestern River. In *AFS 142nd Annual Meeting*. AFS.
- Williams, J.D., M.L. Warren, K.S. Cummings, J.L. Harris & R.J. Neves. 1993. Conservation status of freshwater mussels of the United States and Canada. *Fisheries*. 18(9):6-22.
- Williams, J.D., A.E. Bogan, and J.T. Garner. 2008. *Freshwater Mussels of Alabama & the Mobile Basin in Georgia, Mississippi & Tennessee*. University of Alabama Press: Tuscaloosa, Alabama. 908 pp.
- Young, M. & Williams, J. (1984). The reproductive biology of the freshwater pearl mussel *Margaritifera margaritifera* (Linn.) in Scotland. II Laboratory studies. *Arch. Hydrobiol.* 100, 29–43.
- Yeager, B. (1993). Dams. Pages 57-92 in C.F. Bryan and D. A Rutherford, editors. *Impacts on warm water streams: guidelines for evaluation*. *American Fisheries Society*, Little Rock, Arkansas.